

Product datasheet for **MR211186L4V**

Brd8 (BC025644) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	Brd8 (BC025644) Mouse Tagged ORF Clone Lentiviral Particle
Symbol:	Brd8
Synonyms:	SMAP, 4933408B17Rik
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
Tag:	mGFP
ACCN:	BC025644
ORF Size:	2760 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR211186).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	BC025644 , AAH25644
RefSeq Size:	3277 bp
RefSeq ORF:	2762 bp
Locus ID:	78656
Cytogenetics:	18 B1



[View online »](#)

Gene Summary:

May act as a coactivator during transcriptional activation by hormone-activated nuclear receptors (NR). Stimulates transcriptional activation by AR/DHTR, ESR1/NR3A1, RXRA/NR2B1 and THRB/ERBA2. Component of the NuA4 histone acetyltransferase (HAT) complex which is involved in transcriptional activation of select genes principally by acetylation of nucleosomal histones H4 and H2A. This modification may both alter nucleosome - DNA interactions and promote interaction of the modified histones with other proteins which positively regulate transcription. This complex may be required for the activation of transcriptional programs associated with oncogene and proto-oncogene mediated growth induction, tumor suppressor mediated growth arrest and replicative senescence, apoptosis, and DNA repair. NuA4 may also play a direct role in DNA repair when recruited to sites of DNA damage. Component of a SWR1-like complex that specifically mediates the removal of histone H2A.Z/H2AFZ from the nucleosome.[UniProtKB/Swiss-Prot Function]