

Product datasheet for MR211218L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Diap3 (BC028920) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Diap3 (BC028920) Mouse Tagged ORF Clone Lentiviral Particle

Symbol: Diap3

Synonyms: Dia2, mDia2, p134MDia2

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

 Tag:
 Myc-DDK

 ACCN:
 BC028920

 ORF Size:
 2787 bp

ORF Nucleotide

OTI Disclaimer:

Sequence:

Locus ID:

The ORF insert of this clone is exactly the same as(MR211218).

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

 RefSeq:
 BC028920.1

 RefSeq Size:
 3770 bp

 RefSeq ORF:
 2789 bp

56419

Cytogenetics: 14 E1





Gene Summary:

Actin nucleation and elongation factor required for the assembly of F-actin structures, such as actin cables and stress fibers (PubMed:10678165, PubMed:23558171). Required for cytokinesis, stress fiber formation and transcriptional activation of the serum response factor (PubMed:10678165, PubMed:23558171). Binds to GTP-bound form of Rho and to profilin: acts in a Rho-dependent manner to recruit profilin to the membrane, where it promotes actin polymerization (PubMed:10678165). DFR proteins couple Rho and Src tyrosine kinase during signaling and the regulation of actin dynamics (PubMed:10678165). Also acts as an actin nucleation and elongation factor in the nucleus by promoting nuclear actin polymerization inside the nucleus to drive serum-dependent SRF-MRTFA activity (PubMed:23558171). [UniProtKB/Swiss-Prot Function]