

Product datasheet for **MR211253**

Hdac7 (NM_019572) Mouse Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Hdac7 (NM_019572) Mouse Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	Hdac7
Synonyms:	5830434K02Rik; HD7; HD7a; Hdac7a; mFLJ00062
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)



[View online »](#)

ORF Nucleotide Sequence:

>MR211253 representing NM_019572
 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCC**GCGATCGCC**

ATGCACAGCCCCGGCGGGCTGCCTGCCTCCAGCCAGACACACCAGGCTCTCAGCCCCAACCCATGG
 ACCTGCGGGTGGGCCAGCGGCCACGGTGGAGCCCCACCAGAGCCTGCGCTGCTGACCCTGCAACACCC
 CCAACGCCTGCACCGCCATCTTCTTCTGCGAGGCTTACACCAGCAACAGCGCTCAGCCGAGCCCATGAGG
 CTCTCCATGGACCCACCAATGCCGGAGCTGCAGGGGGACAGCAGGAGCAAGAATTCGGCAACTTCTCA
 ATAAGACAAGAGCAAGCGAAGTCCGTAGCCAGCAGTGTGGTCAAGCAGAAGCTGGCTGAAGTGATCCT
 GAAGAAACAGCAGGCAGCCCTTGAGAGAACAGTCCATCCCAGCAGCCCCAGTATTCCTACAGAACTTT
 GAGCCCTTGGACACAGAGGGTGTGCCCGCTCCGTGCTTAGCAGCTTCTGCCTCCTGTTCCAGCCTGC
 CCACTGAACCCCCGGAACACTTTCCCTTGCCTAAAACAGTGTCTGAACCCAACTGAAGTTGCGCTACAA
 ACCCAAGAAATCCCTGGAGAGACGAAGAATCCCCTGCTCAGGAAGGAGAGTGCCCCGCCAGCCTTCGG
 AGGAGGCCTGCCGAGACCCTTGGAGATTCTCCCCAGTAGTAGCAGCACACCCGCGTCAGGGTGCAGCT
 CCCCTAATGACAGCGAGCATGGCCCTAACCTGCCTAGGCTCAGAGGCTGATGGTGACCGCAGGACCCA
 TTCAACTTTAGGCCCTCGGGTCTGTACTGGGAACCCCATGCTCCCCTTCTCCTGCACCACGGTCTG
 GAGCCAGAGGCTGGGGCACCTTACCCTCTCGCTGCAACCCATTCTCCTGCTGGACCCCTCAGTCTCTC
 ATGCCCACTGTGGACTGTGCCTGGCCTTGGGCCCTTGCCTTCCACTTTGCCAGCCCTTACTGACCAC
 CGAGCGGCTCTCTGGGTGAGCCTCCATCGACCACTTAACCGACCCGCTCAGAGCCCTGCCCCACAG
 GCCACAGCCTCCCTCTGCTGGCCCCCTGCAGCCCCGCCAGGATCGGCTCAAACCTCACGTCCAGCTGA
 TCAAGCCAGCCATCTCCCCTCCCAGAGCCTGCCAAGCCAGTGAGAAGCCCCGATGCCAGACATACC
 CTCGGCTGAGGACCTAGAGACAGATGGTGGGGAGTGGGACCTATGGCGAATGATGGCCTGGAACATAGG
 GAGTCAGGCCGTGGCCTCCTGAGGGCAGAGGCTCCATTTCTCTGCAGCAGCATCAACAGGTGCCACCT
 GGGAGCAGCAGCATCTAGCCGGCGGCTCTCTCAGGGAAGCCCCGGGACTCCGTGCTGATACCTCTGGC
 CCAGGTTGGACACCGGCCCTGTCCAGAACCAGTCTTCCCAGCAGCACCTGTCTCCCTGCTGAGCCCA
 GAGCCACCTGTGACACCAAGTCTCAACAGCTCAGAGACCTGTACAGGGCTGGTCTATGACTCGG
 TGATGCTGAAACACCAATGTTCTGTGGAGACAACAGCAAGCATCCCGAGCATGCAGGCCGATCCAGAG
 CATCTGGTCCCGCTGCAGGAACGGGTCTCCGACCCAGTGTGAGTGTCTCCGAGGCCAAAGGCTTCC
 CTAGAGGAGCTGCAGTCACTCTGAACGGCAGTGTCTCTACGGCACGAACCCACTCAGCCGCC
 TCAAACCTGGATAACGGGAAGCTTACAGGACTCCTGGCACAGCGGACGTTTGTGATGCTACCTGTGGCGG
 GTTGGGGTGCATACTGACACCATCTGGAACGAGCTGCATTCTCCAATGCAGCCCGCTGGGCTGCGGGC
 AGTGTACCGACCTTGCCTTCAAAGTAGCTTCCCAGAGCTGAAGAATGGCTTTGCTGTGGTGGCAGCCC
 CGGGACACCATGCAGATCATTCTACAGCCATGGGCTTCTGCTTCTTCACTCCGTGGCCATCGCCTGCCG
 ACAGCTACAGCAACACGGCAAAGCCAGCAAGATCCTCATTGTTGACTGGGATGTTACCATTGGCAACGGC
 ACACAGCAGACTTCTACCAGGACCCAGTGTGCTCTACATTTCCCTTATCGCCATGACGACGGCAACT
 TCTTCCAGGCAGTGGGGCCGTGGATGAGGTGGAACTGGCAGTGGCGAGGGCTTCAATGTCAACGTGGC
 TTGGGCTGGGGCTTGGATCCACCCATGGGGATCCTGAGTACCTGGCTGCTTTCAGGATAGTGGTGATG
 CCCATTGCCGAGAGTTTGTCCAGACCTGGTCTGTGCTGCTGGGTTTGTGCTGCGGAGGGTACC
 CAGCCCCGCTGGGTGGCTACCATGTTTCTGCCAATGTTTGGGTACATGACGCAGCAGTTGATGAACCT
 GGCAGGAGGCGCGTGGTGTGGCCTTAGAGGGTGGACATGACCTCACGGCCATCTGTGATGCCTCGGAG
 GCCTGTGTAGTGTCTTCTGGGCAACAAGGTGGACCCCTTTCAGAAGAAAGCTGGAAACAGAAACCCA
 ACCTCAGTGCCATCCGCTCGCTGGAAGCTGTGGTCAAGGTGCACAGGAAATACTGGGGCTGCATGCAGCG
 CTTGGCCTCCTGTCCAGACTCCTGGCTACCCAGAGTGCCGGGAGCTGATGCAGAAGTGAAGCCGTGACC
 GCGCTGGCATCCCTTTCTGTGGCATCCTGGCTGAAGACAGGCCCTCGGAGCGGCTGGTGAAGAGGAAG
 AACCCATGAACCTC

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
 ACAAGGATGACGACGATAAGGTTTAA

Protein Sequence: >MR211253 representing NM_019572
 Red=Cloning site Green=Tags(s)


```
MHSPGAGCPALQPDTPGSQPQPMDLRVGQRPTVEPPPEPALLTLQHPQRLHRHLFLAGLHQQRSAEPMR
LSMDPPPELQGGQEQELRQLLNKDKSKRSVAVSSVVKQLAEVILKKQQAALERTVHPSSPSIPYRTL
EPLDTEGAARSVLSFLPPVPSLPTEPEHFPLRKTVSEPNLKLRYKPKKSLERRKNPLLRKESAPPSLR
RRPAETLGDSSPSSSTPASGCSSPNDSEHGPNPALGSEADGDRRTHSTLGPGRPVLGPNHAPLFLHHGL
EPEAGGTLPSRLQPILLLDPSVSHAPLWTVPLGPLPFHFAQPLLTTERLSGSGLHRPLNRTREPLPPS
ATASPLLAPLQPRQDRKPHVQLIKPAISPPQRPAPKSEKPRLRQIPSAEDLETDGGVGPMANDGLEHR
ESGRGPPPEGRGSIISLQQHQVPPWEQQHLAAGRLSQGSPGDSVLIPLAQVGHRLSRTQSSPAAPVLLSP
EPTCQTQVLNSETPATGLVYDSVMLKHQCSCGDNKSHPEHAGRIQSIWSRLQERGLRSQCECLGRKAS
LEELQSVHSERHVLLYGTNPLSRLKLDNGKLTGLLAQRTFVMLPCGGVGVDTDTIWNELHSSNAARWAAG
SVTDLAFKVASRELKNGFAVVRPPGHHADHSTAMGFCFNSVAIACRQLQQHGKASKILIVDWDVHHGNG
TQQTFYQDPSVLVISLHRHDDGNFFPGSGAVDEVTGSGEGFNVNVAWAGGLDPPMGDPEYLAAFRIVVM
PIAREFAPDLVLSAGFDAAEHGPAPLGGYHVSACKCFGYMTQQLMNLAGGAVVLALEGGHDLTAICDASE
ACVAALLGNKVDPLSEESWKQKPNLSAIRSLEAVVRVHRKYWGCMQRLASCPDSWLPRVPGADAEVEAVT
ALASLSVGILAE DRP SERLVEEEEEPMNL
```

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Restriction Sites:

SgfI-MluI

Cloning Scheme:




ACCN: NM_019572

ORF Size: 2814 bp

OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_019572.3
RefSeq Size:	4147 bp
RefSeq ORF:	2817 bp
Locus ID:	56233
UniProt ID:	Q8C2B3
Cytogenetics:	15 F1
MW:	101.7 kDa
Gene Summary:	Responsible for the deacetylation of lysine residues on the N-terminal part of the core histones (H2A, H2B, H3 and H4). Histone deacetylation gives a tag for epigenetic repression and plays an important role in transcriptional regulation, cell cycle progression and developmental events. Histone deacetylases act via the formation of large multiprotein complexes. Involved in muscle maturation by repressing transcription of myocyte enhancer factors such as MEF2A, MEF2B and MEF2C. During muscle differentiation, it shuttles into the cytoplasm, allowing the expression of myocyte enhancer factors. Positively regulates the transcriptional repressor activity of FOXP3 (By similarity). Serves as a corepressor of RARA, causing its deacetylation and inhibition of RARE DNA element binding (By similarity). In association with RARA, plays a role in the repression of microRNA-10a and thereby in the inflammatory response (By similarity).[UniProtKB/Swiss-Prot Function]

Product images:



Circular map for MR211253