

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for MR211329L4V

Kif5b (NM_008448) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	Kif5b (NM_008448) Mouse Tagged ORF Clone Lentiviral Particle
Symbol:	Kif5b
Synonyms:	AL022807; Khc; Khcs; Kns1; Ukhc
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
Tag:	mGFP
ACCN:	NM_008448
ORF Size:	2889 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR211329).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<u>NM 008448.3, NP 032474.2</u>
RefSeq Size:	6030 bp
RefSeq ORF:	2892 bp
Locus ID:	16573
UniProt ID:	<u>Q61768</u>
Cytogenetics:	18 4.46 cM



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Gene Summary:Microtubule-dependent motor required for normal distribution of mitochondria and
lysosomes. May be involved in the mechanisms of growth arrest induced by exposure to
DNA-damaging drugs or by cellular senescence (PubMed:9657148). Can induce formation of
neurite-like membrane protrusions in non-neuronal cells in a ZFYVE27-dependent manner
(PubMed:21976701). Regulates centrosome and nuclear positioning during mitotic entry.
During the G2 phase of the cell cycle in a BICD2-dependent manner, antagonizes dynein
function and drives the separation of nuclei and centrosomes. Required for anterograde
axonal transportation of MAPK8IP3/JIP3 which is essential for MAPK8IP3/JIP3 function in axon
elongation (By similarity).[UniProtKB/Swiss-Prot Function]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US