

Product datasheet for **MR211406L4V**

Epha7 (BC026153) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	Epha7 (BC026153) Mouse Tagged ORF Clone Lentiviral Particle
Symbol:	Epha7
Synonyms:	Ehk3, Mdk1, Cek11, Hek11
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
Tag:	mGFP
ACCN:	BC026153
ORF Size:	2982 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR211406).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	BC026153.1
RefSeq Size:	3554 bp
RefSeq ORF:	2984 bp
Locus ID:	13841
Cytogenetics:	4 12.42 cM



[View online »](#)

Gene Summary:

Receptor tyrosine kinase which binds promiscuously GPI-anchored ephrin-A family ligands residing on adjacent cells, leading to contact-dependent bidirectional signaling into neighboring cells. The signaling pathway downstream of the receptor is referred to as forward signaling while the signaling pathway downstream of the ephrin ligand is referred to as reverse signaling. Among GPI-anchored ephrin-A ligands, EFNA5 is a cognate/functional ligand for EPHA7 and their interaction regulates brain development modulating cell-cell adhesion and repulsion. Has a repellent activity on axons and is for instance involved in the guidance of corticothalamic axons and in the proper topographic mapping of retinal axons to the colliculus. May also regulate brain development through a caspase(CASP3)-dependent proapoptotic activity. Forward signaling may result in activation of components of the ERK signaling pathway including MAP2K1, MAP2K2, MAPK1 AND MAPK3 which are phosphorylated upon activation of EPHA7. Isoform 4 which lacks the kinase domain may regulate isoform 1 adhesive properties.[UniProtKB/Swiss-Prot Function]