

## Product datasheet for MR211479

### Zcchc6 (BC023880) Mouse Tagged ORF Clone

#### Product data:

Product Type:	Expression Plasmids
Product Name:	Zcchc6 (BC023880) Mouse Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	Zcchc6
Synonyms:	MGC38585, 6030448M23Rik
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
ORF Nucleotide Sequence:	>MR211479 representing BC023880 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC  
GCCCGCATCGCC

ATGGCTGTTTTCTTCCTCAGCAAGAAAAGAACCTCTCTTGCCAGTTTATCTAGGATCATGGATTGAAG  
AATTCTCATAAATAAATTAGGGAAATTCAGCCTTAAAGATGTTGAGAAAGACTCTGTGGTCTGGGAATA  
CACTGACAACAGCACAGGGGACACCAGCTCAGCAAAAGAGGAGGCACCCAAAGAGACTGCAGCAAGAAG  
GGACAGGTGCCATTGACATTTAATAAAAGCATCAGCCTTCAGTGCCAGTCGGGCAACTCTGGTGGAAT  
TGCTGCGGTTCTATGCTTTAGAGTTAATCTGGCTGATTTAGTGATCAGTATTCGTGTCAAAGAATTAAT  
ATCACGGGAATCAAAGGACTGGCCAAAAGCGTATCGCAATTGAAGACCCTATTCTGTAAAAAGAAAT  
GTGGCAAGGACCTGAATAACCAGCCTGTGTTGAGTATACTCCATTGTTGAGGACAACATATAAAT  
ATTTGCTCTTCCACACAAAAGTTACAAAACCAACCTCACAAAGCCTCCTAGCCAGTCACCTGTGTTTC  
AGACCCTTACAGAGAAGCAAAAATGGTGGCCAGAACCCCAAGCAACAAATATTGATAAACTGGGAAAT  
GCAGCTGTGGCTCAAGATCCTGGTGTTCAAACCTCAGGTGACTGCAGAGCACAGCTAGTTACTCTTAAGA  
ATACCACTGAGGAGGTGGGAAGCCAGCAAGGAGAAGACAGGAGGTGTGCACATCCCAGCCACCAGGA  
GAGCTCTGGCTGTGTCCAGGCAGAGGTCAGTTGTGAGGGTTTGGAGGATGCCACAGCAGAGCTTCCGGAA  
ACAGGAAGTGACAACGAGGAAGTCAGAAGGAAGACCAAGCATCCTTTGAGCACTGATGACCAAGGTTTAA  
GCAGCAGCAAGCACCCAGAAGTGCAGAAGTGTGGCAGCCTGTGTGGTCTGCAGGCAGACAACACTCTGGA  
GTTAGTAGCTGAAGAGTGAATAGCTGTGCCTCTTTGGACAACAAGGCTGAGGTCAATGAAGAGAGGATA  
GAAGGCCTGAGGAACTAGAAGAAGCAGCAGCTCTGAGTTGCTTTTCCCCTTTCAGTACAGAGTCGCACAT  
CTGCAGCCATGCACTTTGATGATGAGGAGGAGGAGGAGGAAGAAGAGGAGGAAGAGGAGCCAGGCTCAG  
CATTAACTTACAGAAGATGAGGAGGGGTTGCTAATGAGCATCAGGTGGACAGCAGGTATGCTGGGTCA  
GGTGAGGAGGATGCCCTCTGAAGAAGATGACTTGGCAGAGCCTGCTAAGGGTGAAGATACAGGAGAAT  
GTGGAGAAAACGTGGGTGGGACTCTGCTCATAGACCTAAATAGAATAACACTTAAAGAAGAAAGCTTTCC  
TGAGGAAGATTTACCTGGGACCAGTCTGAGTTCTTACGAATTTAGAAAAGTACCTTACCAAAAGGC



[View online »](#)

AAGTCCCCACAGTAGTGTGCAGCTTGTGTAAGCGAGAAGGTCACCTAAAGAAGGACTGTCTGAGGACT  
 TCAAGAGGGTCCAGCTGGAGCCGTTGCCACCCCTGACACCCAAGTTCTCAAACATCTTAGATCAAGTCTG  
 CGTCCAGTGTTACAAGGATTTTTCTCCAATATTGTAGAAGATCAGGCTCGTGAACATATCCGGCAAAAT  
 CTAGAAAGTTTCATAAAAACAGGATTTCCAGGAATAAATTGAGCTTGTGGCTCTCCAAAAATGGAT  
 TTGGGTTCAAACAGAGTGACCTTGACGTCTGTATGACGATTACCGACATGAGACTGTGAGGGGCTGGA  
 CTGTGTAAGAATAATTGAGGAGCTGGCTCGAGTCTCAGAAAGCACTCAGGGTTGAGGAATATCCTTCCT  
 ATCACAACAGCCAAGGTGCCATCGTGAAGTTCTCCATCTAAGAAGTGGTCTGGAAGTGACATCAGCT  
 TGTATAACACACTGGCGCTTATAACACAAGGCTCTGTCTGCATACTCAGCCATTGACCCAGAGTGAA  
 GTATCTGTGCTACACCATGAAAGTGTTCACAAAGATGTGTGACATCGGTGACGCTCCAGAGGCAGCTTG  
 TCATCCTACGCTTACACTCTTATGGTGTATACTTCTCCAGCAAAGGCTCCACCTGTCATCCCCGTGC  
 TTCAAGAGATATACAAAGGTGAAAAGAAACCAGAAATACTTGTGATGGCTGGAATATTTACTTTTTCGA  
 TCAAATAAATGAACTGCCACTTGTGGCCAGAATATGGGAAAAATACAGAACCTGTCCGGCAGCTGTGG  
 CTGGGACTTCTGCGCTTCTACACTGAGGAGTTGACTTTAAGGAGCACGTGATCAGCATCAGAAGAAAGA  
 GTCTGCTTACAACTTTTAAGAAACAGTGGACCTCAAAATACATTGTTATTGAAGATCCCTTCGATTTGAA  
 TCACAATCTTGGTGTGGATTATCGAGGAAAATGACAAATTTTATAATGAAAGCTTTTATCAATGGTAGA  
 AGAGTATTTGGAATCCAGTCAAAGGATTTCAAAGGACAACCCCTCAAAACTGGCATACTTTTTTGATC  
 CGGATGTCCTAACTGAGGGGAACTGGCCCAAATGACAGGTGCTGCCGAATTTGTGGGAAAATCGGGCA  
 CTTTATGAAGGACTGTCCATGAGGAGGAAAGTGAAGGACGAGAGACCAGGAAGATACCCCAAACCAA  
 AGGTACTCTGAGAGCAAAGAAAAAGAAAGCAAAGAGGACAAAGAGATTCAGAACAAGTACACAGAAAAGG  
 AAGTGTCAACAAAAGAAGATAAGCTCACACCGTGTGCAGCTGCCAAAGCCAAGCCAGTGAGGGCAGCTGT  
 CGACCTGGGCCGGGAGAAGCTCCTCAGGACGCCACAGAAAAGTGAAGAGACAGGACGACAGAGACTCG  
 AGAGAAAAGCGTTGTTTCATCTGTGGAAGAGAAGGACACATTAAGGAATGCCACAGTTCAAAGGCT  
 CTCACAGTAGCCTTTCCAGTAAATATATGACTCAGGGGAGAGCCTCAGTGAAGAGGACCCAGCAGGAATC  
 A

AGCGGACCGACGCGTACGCGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCC  
 TGGATTACAAGGATGACGACGATAAGGTTTAA

**Protein Sequence:**


>MR211479 representing BC023880  
 Red=Cloning site Green=Tags(s)

MAVFFLQQRKEPLL PVYLGSWIEEFSLNKLGNFSLKDVEKDSVVWEYTDNSTGDTSSAKEEAPKETAARK  
 GQVPLTFNIKHQPSVPVQQLWVELLRFYALEFNADLVI SIRVKELISRESKDWPKKRIAIEDPYSVKRN  
 VARTLNNQPVFEYILHCLRTTYKYFALPHKVTKPNLTKPPSPVTCVSDPYREAKNGGPEPQATNIDKLG  
 AAVAQDPGVQTS GDCRAQLVTLKNTTEEVGSPAKEKTGGVHIPAHQESSGCVQAEVSCGLEDATAELPE  
 TGS DNEEVRRKTKHPLSTDDQGLSSSKHPELQNCGSLCGLQADNTLELVAEECNASCASLDNKAEVNEERI  
 EGAEEL EEAALSCFSPSVQSR TSAAMHFDDEEEEEEEEEEPRLSINL TEDEEGVANEHQVDSRYAGS  
 GEEDALSEEDLAEPKGEDTGECEGENVGGTLLIDLNRITLKEESFPEEDLPGDQSEFFYEFRLTFTKG  
 KSPTVVCSLCKREGHLKKDCPEDFKRVQLEPLPPLTPKFSNILDQVCVQYKDF SPTIVEDQAREHIRQN  
 LESFIKQDFPGTKLSLFGSSKNGFGFKQSDLDVCMTINGHETAEGLCDVRTIEELARVLRKHSGLRNILP  
 ITTAKVPVIVKFFHLRSGLEVDISLYNTLALHNTRLLSAYS AIDPRVKYL CYTMKVFTKMDIGDASRGS  
 SSYAYTLMLVLYFLQQRSPVIVLQEIYKGEKKPEILVDGWNIIYFFDQINELPTCWPEYGNTEPVGQLW  
 LGLLRFYTEEFDFKEHVISIRRKSLTTFKKQWTSKYI VIEDPFDLNHNLGAGLSRKMTNFIMKAFINGR  
 RVFGIPVKGFPKDNPSKLAYFFDPDVLTEGELAPNDRCCRICGKIGHFMKDCPMRRKVRRRRDQEDTPNQ  
 RYSESKERKSKEDKEIQNKYTEKEVSTKEDKLTPCAA AKAPVRAAVDLGREKLLRTPTEKWKQRQDRDS  
 REKRCFICGREGHIKKECPQFKGSPGSLSSKYMTQGRASVKRTQQES

SGPTRRRLEQKLI SEEDLAANDILDYKDDDDKV

**Restriction Sites:**

SgfI-RsrII

**Cloning Scheme:**


**ACCN:** BC023880

**ORF Size:** 3081 bp

**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

- Reconstitution Method:**
1. Centrifuge at 5,000xg for 5min.
  2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
  3. Close the tube and incubate for 10 minutes at room temperature.
  4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
  5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

**Note:** Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.

**RefSeq:** [BC023880.1](#)

**RefSeq Size:** 4215 bp

RefSeq ORF: 3083 bp


Locus ID: 214290

Cytogenetics: 13 B2

MW: 154.6 kDa

**Gene Summary:** Uridyltransferase that mediates the terminal uridylation of mRNAs with short (less than 25 nucleotides) poly(A) tails, hence facilitating global mRNA decay (PubMed:28792939). Essential for both oocyte maturation and fertility. Through 3' terminal uridylation of mRNA, sculptors, with TUT7, the maternal transcriptome by eliminating transcripts during oocyte growth (PubMed:28792939). Involved in microRNA (miRNA)-induced gene silencing through uridylation of deadenylated miRNA targets. Also acts as a suppressor of miRNA biogenesis by mediating the terminal uridylation of miRNA precursors, including that of let-7 (pre-let-7) (PubMed:22898984). Uridylated pre-let-7 RNA is not processed by Dicer and undergo degradation. Pre-let-7 uridylation is strongly enhanced in the presence of LIN28A. Due to functional redundancy between ZCCHC6 and ZCCHC11, the identification of the specific role of each of these proteins is difficult (By similarity) (PubMed:22898984). Involved in microRNA (miRNA)-induced gene silencing through uridylation of deadenylated miRNA targets (By similarity). Also functions as an integral regulator of microRNA biogenesis using 3 different uridylation mechanisms (By similarity). Acts as a suppressor of miRNA biogenesis by mediating the terminal uridylation of some miRNA precursors, including that of let-7 (pre-let-7). Uridylated pre-let-7 RNA is not processed by Dicer and undergo degradation. Pre-let-7 oligouridylation is strongly enhanced in the presence of LIN28A (PubMed:22898984). In the absence of LIN28A, TUT7 and TUT4 monouridylate group II pre-miRNAs, which includes most of pre-let7 members, that shapes an optimal 3' end overhang for efficient processing (By similarity). Add oligo-U tails to truncated pre-miRNAs with a 5' overhang which may promote rapid degradation of non-functional pre-miRNA species (By similarity). Does not play a role in replication-dependent histone mRNA degradation (By similarity). Due to functional redundancy between TUT4 and TUT7, the identification of the specific role of each of these proteins is difficult (PubMed:28792939, PubMed:22898984). TUT4 and TUT7 restrict retrotransposition of long interspersed element-1 (LINE-1) in cooperation with MOV10 counteracting the RNA chaperone activity of L1RE1. TUT7 uridylates LINE-1 mRNAs in the cytoplasm which inhibits initiation of reverse transcription once in the nucleus, whereas uridylation by TUT4 destabilizes mRNAs in cytoplasmic ribonucleoprotein granules (By similarity).[UniProtKB/Swiss-Prot Function]

Product images:



Circular map for MR211479