

Product datasheet for MR211587

Tek (BC050824) Mouse Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Tek (BC050824) Mouse Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	Tek
Synonyms:	AA517024; Cd202b; Hyk; STK1; Tie-2; Tie2
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
Cell Selection:	Neomycin
ORF Nucleotide Sequence:	>MR211587 representing BC050824 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGGATCGCC**

ATGGACTCTTTAGCCGGCTTAGTTCTCTGTGGAGTCAGCTTGCTCCTTTATGGAGTAGTAGAAGGCGCCA
TGGACCTGATCTTGATCAATCCCTACCTCTTGCTGTGATGCCGAAACATCCCTCACCTGCATTGCCTC
TGGGTGGCACCCCATGAGCCCATCACCATAGGAAGGGACTTTGAAGCCTTAATGAACCAGCACCAAGAT
CCACTGGAGGTTACTCAAGATGTGACCAGAGAATGGGCGAAAAAGTTGTTTGGAAAGAGAGAAAAAGGCCA
GTAAGATTAATGGTGCTTATTTCTGTGAAGGTCGAGTTTCGAGGACAGGCTATAAGGATACGGACCATGAA
GATGCGTCAACAAGCATCCTTCCTACCTGCTACTTTAACTATGACCGTGGACAGGGGAGATAATGTGAAC
ATATCTTTCAAAAAGGTGTTAATTAAGAAGAAGATGCAGTGATTTACAAAAATGGATGTGAAGCTCAGA
AGTGGGGCCCGACTGTAGCCGTCCTTGACTACTTGCAAGAACAATGGAGTCTGCCATGAAGATACCGG
GGAATGCATTTGCCCTCCTGGGTTTATGGGGAGAACATGTGAGAAAGCTTGTGAGCCGCACACATTTGGC
AGGACCTGTAAGAAAAGGTGTAGTGGACCAGAAGGATGCAAGTCTTATGTGTTCTGTCTCCAGACCCTT
ACGGGTGTTCTGTGCCACAGGCTGGAGGGGTTGCAGTGCAATGAAGCATGCCCATCTGGTTACTACGG
ACCAGACTGTAAGCTCAGGTGCCACTGTACCAATGAAGAGATATGTGATCGGTTCCAAGGATGCCTCTGC
TCTCAAGGATGGCAAGGCTGCAGTGTGAGAAAAGGCAAGGCCAAGGATGACTCCACAGATAGAGGATT
TGCCAGATCACATTGAAGTAAACAGTGGAAAATTTAACCCCATCTGCAAAGCCTCTGGGTGGCCACTACC
TACTAGTGAAGAAATGACCCTAGTGAAGCCAGATGGGACAGTGTCCAACCAATGACTTCAACTATACA
GATCGTTTCTCAGTGGCCATATTCAGTGTCAACCGAGTCTTACCTCTGACTCAGGAGTCTGGGTCTGCA
GTGTGAACACAGTGGCTGGGATGGTGGAAAAGCCTTTCAACATTTCCGTCAAAGTTCTTCCAGAGCCCT
GCACGCCCAAAATGTGATTGACTGGACATAACTTTGCTATCATCAATATCAGCTCTGAGCCTTACTTT
GGGGATGGACCCATCAAATCCAAGAAGCTTTTCTATAAAGCTGTCAATCAGGCCTGGAAATACATTGAAG
TGACGAATGAGATTTTCACTCTCAACTACTTGGAGCCCGGACTGACTACGAGCTGTGTGTGAGCTGGC
CCGTCTGGAGAGGGTGGAGAAGGGCATCTGGGCTGTGAGACGATTTACAACAGCGTCTATCGGACTC
CCTCTCCAAGAGGTCTCAGTCTCTGCCAAAAAGCCAGACAGCTCTAAATTTGACTTGGCAACCGATAT



[View online >](#)

TTACAAACTCAGAAGATGAATTTTATGTGGAAGTCGAGAGGCGATCCCTGCAAACAACAAGTGATCAGCA
 GAACATCAAAGTGCCTGGGAACCTGACCTCGGTGCTACTGAGCAACTTAGTCCCCAGGGAGCAGTACACA
 GTCCGAGCTAGAGTCAACACCAAGGCGCAGGGGAGTGGAGTGAAGAAGTCAAGGCTGGACCCTTAGTG
 ACATTCTCCCTCAACAGAAAACATCAAGATCTCCAACATCACTGACTCCACAGCTATGGTTTCTTG
 GACAATAGTGGATGGCTATTCGATTTCTCCATCATCATCCGGTATAAGGTTCAAGGCAAAAATGAAGAC
 CAGCACATTGATGTGAAGATCAAGAATGCTACCGTTACTCAGTACCAGCTCAAGGCTCAGAGCCAGAGA
 CTACATACCATGTGGATATTTTTGCTGAGAACAACATAGGATCAAGCAACCCAGCCTTTTCTCATGAAT
 GAGGACGCTTCCACATTCAGCCTCTGCAGACCTCGGAGGGGAAAGATGCTACTCATAGCCATCCTT
 GGGTCCGCTGGAATGACTTGCATCACCGTGTGTTGGCGTTTCTGATTATGTTGCAACTGAAGAGAGCAA
 ATGTCCAAAGGAGAATGGCTCAGGCATTCCAGAACGTGAGAGAAGAACCAGCTGTGCAGTTAACTCAGG
 AACTCTGGCCCTAACAGGAAGGCCAAAAACAATCCGGATCCCACAATTTATCCTGTGCTGACTGGAAT
 GACATCAAGTTTCAAGACGTGATCGGAGAGGGCAACTTTGGCCAGGTTCTGAAGGCACGCATCAAGAAGG
 ATGGGTTACGGATGGATGCCCCATCAAGAGGATGAAAGAGTATGCCTCAAAGATGATCACAGGGACTT
 CGCAGGAGAACTGGAGGTTCTTTGTAACCTTGACACCATCCAACATCATTAACTCTTTGGGAGCATGT
 GAACACCGAGGCTATTTGTACCTAGCTATTGAGTATGCCCGCATGGAAACCTCTGGACTTCTCGCGTA
 AGAGCAGAGTGTAGAGACAGACCTGCTTTTGCCATCGCCAACAGTACAGCTTCCACACTGTCTCCCA
 ACAGCTTCTTATTTGCTGCAGATGTGGCCCGGGGATGGACTACTTGAGCCAGAAACAGTTTATCCAC
 AGGGACCTGGCTGCCAGAAACATTTTAGTTGGTAAAACCTACATAGCCAAAATAGCAGATTTTGGATTGT
 CACGAGGTCAAGAAGTGTATGTAAAAAGACAATGGGAAGGCTCCCAGTGCCTGGATGGCAATCGAATC
 ACTGAACATAGTGTCTATAACAACAACAGTGTGTCTGGTCTATGGTGTATTGCTCTGGGAGATTGTT
 AGCTTAGGAGGCCCCCTACTGCGGCATGACGTGCGCGGAGCTCTATGAGAAGTACCCAGGGCTACA
 GGCTGGAGAAGCCCTGAACTGTGATGATGAGGTGTATGATCTAATGAGACAGTGTGGAGGGAGAAGCC
 TTATGAGAGACCATCATTTGCCAGATATTGGTGCCTTAAACAGGATGCTGGAAGAACGGAAGACATAC
 GTGAACACCACACTGTATGAGAAGTTTACCTATGCAGGAATTGACTGCTCTGCGGAAGAAGCAGCC

ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAATGATATCCTGGATT
 ACAAGGATGACGACGATAAGGTTTAA

Protein Sequence:

>MR211587 representing BC050824
 Red=Cloning site Green=Tags(s)

MDSL AGLVLCGVSLLLYGVVEGAMDILINSLPLVSDAETSLTCIASGWHPHEPITIGRDFEALMNQHQD
 PLEVTDVDTREWAKKVVWKREKASKINGAYFCEGRVVRGQAIRIRTMKMRQASFLPATLTMTVDRGDNVN
 ISFKKVLIKEEAVIYKNGCEAQKWGPDCSRPCTTCKNNGVCHEDTGEICPPGFMGRTCEKACEPHTFG
 RTCKERCSGPEGCKSYVFLPDPYGCSCATGWRGLQCNEACPSGYYGPDCKLRCHCTNEEICDRFQGCLC
 SQGWQGLQCEKEGRPRMTPQIEDLPDHIEVNSGKFNPIKASGWPLPTSEEMTLVKPDGTVLQPNDFNYT
 DRFSVAIFTVNRVLPDPSGVVWCSVNTVAGMVEKPFNISVKVLEPLHAPNVIDTGHNFAIINISSEPYF
 GDGPIKSKKLFYKPVNQAWKYIEVTNEIFTLNYLEPRTDYELCVQLARPEGEGGHPGPVRRFTTASIGL
 PPRGLSLLPKSQTALNLWQPIFTNSEDEFYVEVERRSLQTTSDQQNIKVPGNLTSVLLSNLVPREQYT
 VRRARVNTKAQGEWSEELRAWLSDILPPQENIKISNITDSTAMVSWTIVDGYSISSIIIRYKVGKQKNE
 QHIDVKIKNATVTQYQLKLEPETTYHVDIFAENNISSNPAFSHELRTLPHSPASADLGGGKMLLIAIL
 GSAGMTCITVLLAFLIMLQLKRANVQRRAQAFQNVREEPAVQFNSGTLALNRKAKNPDPTIYPVLDWN
 DIKFQDVI GEGNFQVLKARIKKDGLRMDAAIKRMKEYASKDDHRDFAGELEVLCCKLGHHPNIIINLLGAC
 EHRGYLYLAIEYAPHGNLLDFLRKSRVLETDPAFAIANSTASTLSSQQLLHFAADVARGMDYLSQKQFIH
 RDLAARNILVGENYIAKIADFGLSRGQEVYVKKTMGRLPVRWMAIESLNYSVYTTNSDVWSYGVLLWEIV
 SLGGTPYCGMTCAELYEKLPQGYRLEKPLNCDDEVYDLMRQCWREKPYERPSFAQILVSLNRMLEERKTY
 VNTTLYEKFTYAGIDCSAEAAA

TRTRPLEQKLI SEEDLAANDILDYKDDDDKV


Restriction Sites:

Sgfl-MluI

Cloning Scheme:



Plasmid Map:



ACCN: BC050824

ORF Size: 3216 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	BC050824.1
RefSeq Size:	4549 bp
RefSeq ORF:	3218 bp
Locus ID:	21687
Cytogenetics:	4 43.34 cM
MW:	166.8 kDa
Gene Summary:	<p>Tyrosine-protein kinase that acts as cell-surface receptor for ANGPT1, ANGPT2 and ANGPT4 and regulates angiogenesis, endothelial cell survival, proliferation, migration, adhesion and cell spreading, reorganization of the actin cytoskeleton, but also maintenance of vascular quiescence. Has anti-inflammatory effects by preventing the leakage of proinflammatory plasma proteins and leukocytes from blood vessels. Required for normal angiogenesis and heart development during embryogenesis. Required for post-natal hematopoiesis. After birth, activates or inhibits angiogenesis, depending on the context. Inhibits angiogenesis and promotes vascular stability in quiescent vessels, where endothelial cells have tight contacts. In quiescent vessels, ANGPT1 oligomers recruit TEK to cell-cell contacts, forming complexes with TEK molecules from adjoining cells, and this leads to preferential activation of phosphatidylinositol 3-kinase and the AKT1 signaling cascades. In migrating endothelial cells that lack cell-cell adhesions, ANGPT1 recruits TEK to contacts with the extracellular matrix, leading to the formation of focal adhesion complexes, activation of PTK2/FAK and of the downstream kinases MAPK1/ERK2 and MAPK3/ERK1, and ultimately to the stimulation of sprouting angiogenesis. ANGPT1 signaling triggers receptor dimerization and autophosphorylation at specific tyrosine residues that then serve as binding sites for scaffold proteins and effectors. Signaling is modulated by ANGPT2 that has lower affinity for TEK, can promote TEK autophosphorylation in the absence of ANGPT1, but inhibits ANGPT1-mediated signaling by competing for the same binding site. Signaling is also modulated by formation of heterodimers with TIE1, and by proteolytic processing that gives rise to a soluble TEK extracellular domain. The soluble extracellular domain modulates signaling by functioning as decoy receptor for angiopoietins. TEK phosphorylates DOK2, GRB7, GRB14, PIK3R1, SHC1 and TIE1.[UniProtKB/Swiss-Prot Function]</p>