

Product datasheet for MR211638

Pik3cg (NM_001146201) Mouse Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Pik3cg (NM_001146201) Mouse Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	Pik3cg
Synonyms:	5830428L06Rik; p110gamma; PI3Kgamma
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
ORF Nucleotide Sequence:	>MR211638 representing NM_001146201 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGATCGCC**

ATGGAGCTGGAGAACTATGAACAACCGGTGGTTCTAAGAGAGGACAACCTCCGCCGGCGCCGAGGATGA
AGCCACGCAGCGCAGCAGGCAGCCTGTCTTCCATGGAGCTCATCCCCATTGAGTTCGACTGCCACCAG
CCAGCGCATCAGCAAGACTCCAGAAACAGCGCTGCTGCATGTGGCTGGCCATGGCAATGTGGAACAGATG
AAAGCTCAGGTGTGGCTGCGCGCACTGGAGACCAGTGTGGCTGCGGAGTCTACCACCGATTGGGCCGG
ACCAATTCCTCCTGCTCTACCAGAAGAAAGGACAATGGTATGAGATCTATGACAGGTACCAAGTGGTGCA
GACCCTAGACTGCCTGCATTACTGGAAGTTGATGCACAAGAGCCCTGGCCAGATCCACGTGTACAGCGA
CACGTACCTTCTGAGGAGACCTTGGCTTCCAGAAGCAGCTCACCTCCCTGATTGGCTATGACGTCACTG
ACATCAGCAATGTGCACGATGATGAGCTAGAGTCACTCGCCCGCTGTGGTTACGCCCCGCATGGCTGA
AGTGGCTGGCCGGGATGCCAAACTCTATGCTATGCACCCTTGGGTAACGTCCAAACCTCTCCAGACTAC
CTGTCAAAAAGATTGCCAAACTGCATCTTCATCGTCATCCACCGCGTACCACCAGCCAAACCATCA
AGGTCTCCGCAGATGATACTCCTGGTACCATCCTCCAGAGCTTCTCACCAAGATGGCCAAGAAGAAGTC
CCTAATGAATATCTCAGAAAGTCAAAGTGAGCAGGATTTTGTATTGCGGGTTTGTGGCCCGATGAGTAC
CTGGTGGGTGAAACACCCCTCAAAAATTTCCAGTGGGTGAGGCAGTGCCCAAGAACGGAGATGAAATAC
ACCTGGTGTCTCGACACCGCTCCAGACCCAGCCCTTGTGAGGTGAGGAAGGAAGAATGGCCGCTGGTGGA
TGA CTGCACTGGAGTCAACCGCTACCACGAGCAGCTGACCATCCATGGCAAGACCACGAGAGTGTGTTT
ACAGTGTCTTTGTGGACTGCGACCGAAAGTTCAGGGTCAAGATCAGAGGCATTGATATCCCTGTCTGCTGC
CTCGGAACACCGACCTCACTGTGTTTGTGGAAGCGAACATCCAGCACGGGCAACAAGTCTCTGCCAAAG
GAGAACCAGCCCTAAGCCCTTCGAGAAGAGGTA CTCTGGAATGTGTGGCTGGAGTTTGGCATCAAATC
AAAGACTTGCCCAAGGGGCTCTATTGAACCTACAGATCTACTGCTGCAAAACCCATCACTGTCCAGCA
AGGCTTCTGCAGAGACTCCAGGCTCCGAGTCCAAGGGCAAGCCAGCTTCTCTATTACGTGAACCTTGT
GTTAATAGACCACCGTTTCTCCTCCGCCAGGGGACTATGTGCTCCACATGTGGCAGATATCTGGCAAG



[View online »](#)

GCAGAGGAGCAGGGCAGCTTCAATGCTGACAAGCTCACATCCGCAACCAATCCTGACAAGGAGAACTCAA
 TGTCCATTTCCATCCTGCTGGACAATTACTGTACCCCATAGCTTTGCCTAAGCACCGGCCACCCCTGA
 CCCAGAGGGAGACAGGGTTGGGCTGAAATGCCAATCAGCTTCGAAAGCAATTGGAGGCGATCATAGCC
 ACAGATCCACTTAACCCCTCACAGCAGAGGACAAAGAATTGCTCTGGCATTTCGATATGAAAGCCTGA
 AGCATCCGAAGGCTTACCCTAAGCTATTCAGCTCAGTAAATGGGGCAGCAAGAATTGTTGCCAAAAC
 GTACCAGCTGTTAGCCAGAAGGGAGATCTGGGATCAAAGTGCTTTGGACGTTGGCTTAACCATCAGAGCTC
 CTGGACTGCAACTTTTCAGACGAGAATGTCCGGGCCATTGCAGTTCAGAAACTGGAGCTTAGAGGACG
 ATGACGTTTTACATTACCTTCTCCAGCTGGTACAGGCTGTGAAATTTGAACCGTACCACGACAGTGCCT
 GGCCAGATTCTGCTGAAGCGTGGCTTGAAGAACAAAAGAATCGTCACTTCTTGTCTGGTTCTGCGA
 AGTGAGATCGCACAGTCCAGACACTATCAGCAGAGGTTGCTGTGATCCTGGAGGCGTACCTGCGAGGCT
 GTGGCAGACCCATGTTGCAGGACTTCACACAGCAGGTCCATGTGATTGAGATGTTACAGAAAGTCACCAT
 TGATATTAATCGCTCTCGGCAGAGAAGTATGACGTGAGTCCCAAGTTATTTACAGCTTAAGCAAAG
 CTTGAAAGCCTTCAGAACTCCAATCTCCCCGAGAGCTTTAGAGTTCCTATGATCCTGGACTAAAAGCCG
 GTACCCTGGTATCGAGAAATGCAAAGTATGGCCTCCAAGAAGAAGCCCTGTGGCTTGAGTTAAGTG
 TGCTGATCCCACAGTCTATCCAACGAAACCATTGGAATCATCTTTAAACATGGTATGATCTGCGCCAA
 GACATGTTGATCTTGAGATTCTACGCATCATGGAGTCCATTTGGGAGACTGAATCTCTGGACCTGTGCC
 TTCTGCCTTACGGTTGCATCTCAACTGGTGACAAAATAGGAATGATCGAGATTGTAAGGATGCCACAAC
 GATCGCTCAAATTCAGCAAAGCACAGTGGGTAACACGGGGCATTCAAAGATGAAGTCTGAATCACTGG
 CTCAAGGAAAAATGTCCTATTGAAGAAAAGTTTTCAGGCCGAGTGGAAAGGTTTGTCTTACTCCTGTGCG
 GCTACTGTGTGGCCACATTTGTTCTTGGGATCGGTGACAGGCACAACGACAACATTATGATCTCAGAGAC
 AGGAAACCTATTTCATATAGACTTCGGACACATTCTTGGGAATACAAGAGTTTCTGGGCATCAATAAA
 GAGAGAGTGCCCTTCGTCTAACCCAGACTTCTTGTGTTGATGGGATCTTCTGAAAAAAGACAAGTC
 CACACTCCAGAAATCCAGGATGTCTGTGTTAGAGCTTACCTAGCTCTTCCGCATCACACAAACTGTT
 GATCATCTTGTCTCCATGATGCTGATGACAGGAATGCCCCAGCTGACAAGCAAAGAGGACATTGAATAT
 ATCCGGGATGCCCTACCGTGGGAAAAAGCAGGAGGACGCTAAGAAATATTTCTTGTGATCAGATCGAAG
 TCTGCAGAGACAAAGGATGGACTGTGCAGTTTAACTGGTCTACATCTTGTCTTGGCATCAAACAAGG
 AGAAAAGCACTCCGCT

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
 ACAAGGATGACGACGATAAGGTTTAA

Protein Sequence:

>MR211638 representing NM_001146201
 Red=Cloning site Green=Tags(s)

MELENYEQPVVLRDNLRRRRRMRPRSAAGSLSSMELIPIEFVLPTSQRISKTPETALLHVAGHGNVEQM
 KAQVWLRALETSVAAEFYHRLGPDQFLLLYQKKGQWYIYDRYQVVQTLDCLEHYWKLMMHSPGQIHVVQR
 HVPSEETLAFQKQLTSLIGYDVTDISNVHDELEFTRRRLVTPRMAEVAGRDAKLAMHPWVTSKPLPDY
 LSKKIANNCIFIVIHRTTSQTIKVSADDTPGTILQSFFTKMAKKSLMNISESQSEQDFVLRVCGRDEY
 LVGETPLKNFQWVRQCLKNGDEIHLVLDTPDPALDEVKKEWPLVDDCTGVTGYHEQLTIHGKDHESVF
 TVSLWDCDRKFRVKIRGIDIPVLPNTDLTVFVEANIQHGGQVLCQRRTPSPKPFEEVWVWLEFGIKI
 KDLPKGALLNLQIYCKTPSLSSKASAETPGSESKGAQLLYVYNLLIDHRFLLRHGDYVLLHMWQISGK
 AEEQGSFNADKLTSATNPDKENSMSISILLDNYCHPIALPKHRTPDPEGDRVRAEMPNLQRKQLEAIIA
 TDPLNPLTAEDKELLWHFRYELKHPKAYPKLFSSVKWQQEIVAKTYQLLARREIWDQSALDVGLTMQL
 LDCNFDENVRAIAVQKLESLEDDDLHYLLQLVQAVKFEPYHDSALARFLLKRGRLRNKRIGHFLFWFLR
 SEIAQSRHYQRFVAVILEAYLRGCGTAMLQDFTQQVHVIEMLQKVTIDIKSLSAEKYDVSSQVISQLKQK
 LESLQNSNLPESFRVPYDPGLKAGTLVIEKCKVMASKKPLWLEFKCADPTVLSNETIGIIFKHGDDLQ
 DMLILQILRIMESIWETESLDLCLLPYGCISTGDKIGMIEIVKDATTIAQIQSTVGNLTGAFKDEVLNHW
 LKEKCPIEEFQAAVERFVYSCAGYCVATFVLGIGDRHNDNIMISSETGNLFHIDFGHILGNYSFLGINK
 ERVPFVLPDFLFVMSGSKKTSPLHFQKFDVQVRAVYLRHHTNLLIILFSMMLMTGMPQLTSKEDIEY
 IRDALTVGKSEEDAKKYFLDQIEVCRDKGWTQVFNWFLHLVLGIKQGEKHS

TRTRPLEQKLI SEEDLAANDILDYKDDDDKV

Restriction Sites: SgfI-MluI

Cloning Scheme:



ACCN: NM_001146201

ORF Size: 3306 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_001146201.2](#)

RefSeq Size: 6690 bp

RefSeq ORF: 3309 bp

Locus ID: 30955


UniProt ID: [Q9JHG7](#)

Cytogenetics: 12 A3

MW: 126.8 kDa

Gene Summary: Phosphoinositide-3-kinase (PI3K) that phosphorylates PtdIns(4,5)P₂ (Phosphatidylinositol 4,5-bisphosphate) to generate phosphatidylinositol 3,4,5-trisphosphate (PIP₃). PIP₃ plays a key role by recruiting PH domain-containing proteins to the membrane, including AKT1 and PDK1, activating signaling cascades involved in cell growth, survival, proliferation, motility and morphology. Links G-protein coupled receptor activation to PIP₃ production. Involved in immune, inflammatory and allergic responses. Modulates leukocyte chemotaxis to inflammatory sites and in response to chemoattractant agents. May control leukocyte polarization and migration by regulating the spatial accumulation of PIP₃ and by regulating the organization of F-actin formation and integrin-based adhesion at the leading edge. Controls motility of dendritic cells. Together with PIK3CD is involved in natural killer (NK) cell development and migration towards the sites of inflammation. Participates in T-lymphocyte migration. Regulates T-lymphocyte proliferation and cytokine production. Together with PIK3CD participates in T-lymphocyte development. Required for B-lymphocyte development and signaling. Together with PIK3CD participates in neutrophil respiratory burst. Together with PIK3CD is involved in neutrophil chemotaxis and extravasation. Together with PIK3CB promotes platelet aggregation and thrombosis. Regulates alpha-IIb/beta-3 integrins (ITGA2B/ITGB3) adhesive function in platelets downstream of P2Y₁₂ through a lipid kinase activity-independent mechanism. May have also a lipid kinase activity-dependent function in platelet aggregation. Involved in endothelial progenitor cell migration. Negative regulator of cardiac contractility. Modulates cardiac contractility by anchoring protein kinase A (PKA) and PDE3B activation, reducing cAMP levels. Regulates cardiac contractility also by promoting beta-adrenergic receptor internalization by binding to GRK2 and by non-muscle tropomyosin phosphorylation. Also has serine/threonine protein kinase activity: both lipid and protein kinase activities are required for beta-adrenergic receptor endocytosis. May also have a scaffolding role in modulating cardiac contractility. Contribute to cardiac hypertrophy under pathological stress. Through simultaneous binding of PDE3B to RAPGEF3 and PIK3R6 is assembled in a signaling complex in which the PI3K gamma complex is activated by RAPGEF3 and which is involved in angiogenesis (By similarity).[UniProtKB/Swiss-Prot Function]

Product images:



Circular map for MR211638