

Product datasheet for MR211651

Upf1 (NM_030680) Mouse Tagged ORF Clone

Product data:

Product Type: Expression Plasmids
Product Name: Upf1 (NM_030680) Mouse Tagged ORF Clone
Tag: Myc-DDK
Symbol: Upf1
Synonyms: B430202H16Rik; NORF1; PNORF-1; Rent1; Upflp
Vector: pCMV6-Entry (PS100001)
E. coli Selection: Kanamycin (25 ug/mL)
Cell Selection: Neomycin
ORF Nucleotide Sequence: >MR211651 representing NM_030680
 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCC**CGATCGCC**

ATGAGTGTGGAGGCGTACGGCCCCAGCTCGAAACACTCACCTTCTTGGACTGAGGAGGCCGAGCTGC
 TCGGCCCGACACCCAGGGCTCCGAGTTCGAATTCACCGACTTACCCTTCCCAGCCAGACGCAGACGCC
 CCCCAGGCGCCCGGGCGCGGGAGGCCCGGGCGGAGCGGGCGCAGGCGCGCGGCCAGCTCGAC
 GCACAAGTTGGACCAGAGGGCATCTTGCAAAATGGGGCTGTGGATGACAGTGTGGCCAAGACCAGCCAGC
 TGCTAGCTGAGCTGAACCTCGAGGAAGATGAAGAGGACACATACTACACTAAGGACCTCCAGTCCACGC
 CTGCAGTTACTGTGGAATCCATGATCCTGCCTGCGTGGTTTACTGTAATACCAGCAAGAAGTGGTTCTGC
 AATGGCCGAGGAAATACTTCTGGCAGCCACATTGTGAATCACCTCGTGAGGGCAAATGCAAGGAAGTGA
 CGCTGCACAAGGACGGCCCTCTGGGCGAGACCGTGTGGAGTGTACAACCTGTGGCTGCCCAACGTCTT
 CCTGCTGGGCTTCATCCCTGCGAAGGCCGACTCTGTGGTGGTGTGTTGTGCAGGCAGCCCTGTGCCAGC
 CAGAGCAGCCTGAAGGACATCAACTGGGACAGCTCACAGTGGCAGCCCTAATCCAGGACCGTGTCTTTC
 TGTCATGGCTGGTCAAGATTCCCTCTGAGCAGGAGCAGCTGCGAGCACGGCAGATCACGGCACAGCAGAT
 CAACAAGCTGGAAGAGCTCTGGAAGGAAAATCCTTCAGCCACTCTGGAGGACCTGGAGAAGCCAGGCGTA
 GACGAGGAGCCACAGCAGCTGCTCCTGCGTTACGAGGATGCTTACCAGTACCAGAACATCTTCGGGCCAC
 TGGTCAAGCTGGAGGCTGACTATGACAAGAAGTTGAAGGAGTACAGACTCAAGATAACATCACGGTCAG
 GTGGGACCTGGGCCTTAACAAGAAGAGAATCGCCTTCTTCACTTTGCCAAGACTGACTCTGACATGCGG
 CTCATGCAGGGTATGAGATCTGTCTGCGGTACAAAGGGGATCTGGCGCCCTGTGGAAGGGGATTGGCC
 ACGTCATCAAGGTTCTGATAATTATGGTATGAGATTGCTATTGAGCTCCGAGCAGCGTGGGTGCCCC
 TGTGGAAGTGACCCACAACCTCCAAGTGGATTTGTGTGGAAGTCAACCTCTTTTATAGGATGCAGAGT
 GCACTGAAGACCTTCGCTGTGGACGAGACCTCTGTGTCAGGGTATATTTACCACAAGCTGCTGGGCCAGC
 AGGTGGAGGATGTGGTCAAGTGCCAGCTGCCAAAGCGCTTACAGCTCAGGGGCTCCCTGACCTCAA
 CCACTCTCAGGTGATGCTGTGAAGACCGTGTGTCAGAGACCACTCAGCCTCATCCAGGGCCCTCCAGGC
 ACAGGCAAGACTGTGACATCAGCCACTATTGTCTACCACCTTGTCTCGGCAGGGCAATGGGCTGTACTGG



TTTGTGCTCCAAGTAACATCGCTGTGGACCAGCTCACAGAGAAGATCCACCAGACAGGACTGAAGGTCTG
 ACGCCTCTGTGCCAAGAGCCGTGAGGCCATTGACTCCCCAGTGTCTTCTCGGCTTTGCACAACCAGATC
 AGGAACATGGACAGCATGCCTGAGCTGCAGAAGCTGCAGCAGCTAAAGGATGAGACAGGCGAGCTGCAT
 CTGCAGATGAGAAAGCGGTACCGGGCGCTTAAGCGCACAGCTGAGAGAGAAGTCTCATGAATGCAGATGT
 CATATGCTGCACATGTGTGGTGCTGGTGACCCGAGGCTGGCCAAGATGCAGTTCGGTCCATCCTCATT
 GACGAGAGCACCCAGGCCACTGAGCCTGAGTGCATGGTGCCTGTAGTCTTGGGGCAAGCAGCTAATCC
 TCGTCGGTGACCACTGCCAGCTGGGCCAGTGGTGTGATGTGCAAGAAGGCAGCCAAGGCCGACTGCACA
 ATCGCTCTTCGAGCGCTTGGTGGTGTGGGCATCCGGCCATCCGCTGCAGGTGCAATACCGCATGCAC
 CCTGCACTCAGCGCCTTCCGTCCAACATCTTCTACGAGGGCTCATTGCAGAATGGCGTCACTGCAGCGG
 ATCGTGTCAAAAAAGGCTTTGACTTCCAGTGGCCACAACCTGACAAGCCTATGTTCTTCTACGTGACGCA
 GGGCCAGGAGGATTGCCAGCTCTGGCACATCTACCTCAACAGGACGGAGGCAGCCAATGTGGAGAAG
 ATAACACGAAGCTGTTGAAGGCAGGTGCAAGCCTGACCAGATCGGCATCATCACCCCCACGAGGGCC
 AGCGCTTACTTGGTGCAGTACATGCAGTTCAGCGGCTCCCTGCACACAAAGCTCTACCAGGAAGTGA
 AATTGCCAGTGTGACGCCTCCAGGGCCGGGAGAAGGACTTCATCATTCTGTCTCGCTGCGCGCCAAT
 GAACATCAGGGCATTGGGTTCTAAACGACCCCGGCGTCTGAATGTGGCTCTACCAGAGCAAGATATG
 GCGTGATCATTGTGGGTAAACCAAAGGCCCTGTGGAAGCAGCCCTGTGGAATCAGCTGTGAGCTACTA
 CAAGGAACAGAAGGCGCTAGTGAAGGGCCGCTCAACAACCTACGTGAGAGCCTCATGCAGTTCAGCAAG
 CCTCGAAACTTGTCAACACTGTCAACCCGGGTGCCCGCTTATGACTACTGCCATGTACGATGCCCGTG
 AGGCCATCATCCCGGCTGTCTATGACCGCAGCAGCCAGGGCCGGCCCTCGAACATGTACTTCCAGAC
 CCATGACCAGATTAGTATGATCAGCGCAGGCCAGCCACGTGGCTGCCATGAACATCCCTATTCCCTTC
 AACTTGGTCATGCCTCCCATGCCCCACCTGGCTACTTCGGACAGGCCAACGGGCCGGCAGCTGGTGGG
 GCACCCCAAAAACCAAGACTGGCCGTGGGGCCGCCAGAAGAACCCTTGGGCTTCTGGGCCAGCCA
 GACCACCTTCCCAACAGCCAGGCCAGCAGGACGTGGCTCCAGCCCTTTTACAGGGTGCCTCACA
 CAGGGTTACGTGTCCATGAGCCAGCCCTCTCAGATGAGCCAGCCTGGCCCTCCAGCAGAACTGTCCTC
 AGGACAGTACCTCGGTGATGAGTTAAATCACAGATTGACGTGGCACTCTCACAAGACTCCACATACCA
 GGGAGAGCGGGCATACCAGCACGGCGGGTACCAGGGCTGTCCAGTAC

ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
 ACAAGGATGACGACGATAAGGTTTAA

Protein Sequence:

>MR211651 representing NM_030680
 Red=Cloning site Green=Tags(s)

MSVEAYGPSSQTLTFLDTEEAELLGADTQGSEFEFTDFTLPSQTQTPPGGPGGAGGPGGAGAGGAAGQLD
 AQVGPGLQNGAVDDSAKTSQLLAELNFEDEEDTYTKDLPVHACS YCGIHPACVVCNTSKKWF
 NGRGNTSGSHIVNHLVRAKCKEVLHKDGPLGETVLECYNCGCRNVLLGFIPAKADSVVLLCRQPCAS
 QSSLKDINWDSSQWQPLIQDRCFLSWLVKIPSEQELRARQITAAQQINKLEELWKENPSATLEDLEKPGV
 DEEPQHVLRLRYEDAYQYQNIIFGPLVKLEADYDKLKESTQDNITVRWDLGLNKKRIAFFTLPKTDSMR
 LMQGDEICLRKYGDLAPLWKGIGHVIKVPDNYGDEIAIELRSSVGPVEVTHNFQVDFVWVSTSFDRMQS
 ALKTFVAVDETSVSGYIYHKLLGHEVEDVVIKQQLPKRFTAQGLPDLNHSQYVAVKTVLQRPVSLIQGPPG
 TGKTVTSATIVYHLARQNGPVLVCAPSNIIVDQVTEKIHQTGLKVVRLCAKSREAIIDSPVSLALHNQI
 RNMDMPPELQKLQQLKDETGELSSADEKRYRALKRTAERELLMNADVICCTCVGADPRLAKMQFRSIL
 DESTQATEPECMVPVVLGAKQLILVGDHCQLGPVVMCKKAAKAGLSQSLFERLVVLRIRLQVQYRMH
 PALSAFPSNIFYEGSLQNGVTAADRVKKGFDQWQPDPKPMFFVYVYVQGEIASSGTSYLNRTAANVEK
 ITTKLLKAGAKPDQIGIITPYEQRSYLQYMQFSGSLHTKLYQEVEIASVDAFQGREKDFIILSCVRAN
 EHQIGFLNDPRRLNVALTRARYGVIIIVGNPKALSKQPLWNHLLSYYKEQKALVEGPLNNLRESLMQFSK
 PRKLVNTVNPGRFMTTAMYDAREAIIPGSVYDRSSQGRPSNMYFQTHDQISMISAGPSHVAAMNIPF
 NLVMPPMPPPGYFGQANGPAAGRTPKTKTGRGGRQKNRFGLPGPSQTTLPNQSASQDVASQPF SQALT
 QGYVMSQPSQMSQPLSQPELSQDSYLGDEFKSIDVALSQDSTYQGERAYQHGGVTGLSQY

TRTRPLEQKLISEEDLAANDILDYKDDDDKV


Restriction Sites:

Sgfl-MluI

Cloning Scheme:



Plasmid Map:



ACCN: NM_030680

ORF Size: 3339 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	<u>NM_030680.3</u>
RefSeq Size:	4598 bp
RefSeq ORF:	3342 bp
Locus ID:	19704
UniProt ID:	<u>Q9EPU0</u>
Cytogenetics:	8 34.15 cM
MW:	123.1 kDa
Gene Summary:	<p>RNA-dependent helicase and ATPase required for nonsense-mediated decay (NMD) of mRNAs containing premature stop codons. Is recruited to mRNAs upon translation termination and undergoes a cycle of phosphorylation and dephosphorylation; its phosphorylation appears to be a key step in NMD. Recruited by release factors to stalled ribosomes together with the SMG1C protein kinase complex to form the transient SURF (SMG1-UPF1-eRF1-eRF3) complex. In EJC-dependent NMD, the SURF complex associates with the exon junction complex (EJC) (located 50-55 or more nucleotides downstream from the termination codon) through UPF2 and allows the formation of an UPF1-UPF2-UPF3 surveillance complex which is believed to activate NMD. Phosphorylated UPF1 is recognized by EST1B/SMG5, SMG6 and SMG7 which are thought to provide a link to the mRNA degradation machinery involving exonucleolytic and endonucleolytic pathways, and to serve as adapters to protein phosphatase 2A (PP2A), thereby triggering UPF1 dephosphorylation. UPF1 can also activate NMD without UPF2 or UPF3, and in the absence of the NMD-enhancing downstream EJC indicative for alternative NMD pathways. Plays a role in replication-dependent histone mRNA degradation at the end of phase S; the function is independent of UPF2. For the recognition of premature termination codons (PTC) and initiation of NMD a competitive interaction between UPF1 and PABPC1 with the ribosome-bound release factors is proposed. The ATPase activity of UPF1 is required for disassembly of mRNPs undergoing NMD (By similarity). Essential for embryonic viability.[UniProtKB/Swiss-Prot Function]</p>