

## Product datasheet for MR211687L3

### Nckap1l (NM\_153505) Mouse Tagged Lenti ORF Clone

#### Product data:

Product Type:	Expression Plasmids
Product Name:	Nckap1l (NM_153505) Mouse Tagged Lenti ORF Clone
Tag:	Myc-DDK
Symbol:	Nckap1l
Synonyms:	4930568P13Rik; AI463083; Hem1; Hemp1
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
E. coli Selection:	Chloramphenicol (34 ug/mL)
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR211687).
Restriction Sites:	SgfI-MluI
Cloning Scheme:	

Cloning sites used for ORF Shuttling:



\* The last codon before the Stop codon of the ORF.

ACCN:	NM_153505
ORF Size:	3402 bp



[View online »](#)

<b>OTI Disclaimer:</b>	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a>
<b>OTI Annotation:</b>	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
<b>Components:</b>	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
<b>Reconstitution Method:</b>	<ol style="list-style-type: none"><li>1. Centrifuge at 5,000xg for 5min.</li><li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li><li>3. Close the tube and incubate for 10 minutes at room temperature.</li><li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li><li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li></ol>
<b>RefSeq:</b>	<a href="#">NM_153505.4</a> , <a href="#">NP_705725.1</a>
<b>RefSeq Size:</b>	4717 bp
<b>RefSeq ORF:</b>	3405 bp
<b>Locus ID:</b>	105855
<b>UniProt ID:</b>	<a href="#">Q8K1X4</a>
<b>Cytogenetics:</b>	15 F3
<b>Gene Summary:</b>	Essential hematopoietic-specific regulator of the actin cytoskeleton. Controls lymphocyte development, activation, proliferation and homeostasis, erythrocyte membrane stability, as well as phagocytosis and migration by neutrophils and macrophages (PubMed:19015308, PubMed:23424621). Component of the WAVE2 complex which signals downstream of RAC to stimulate F-actin polymerization (PubMed:23424621). Required for stabilization and/or translation of the WAVE2 complex proteins in hematopoietic cells (PubMed:19015308). Exhibits complex cycles of activation and inhibition to generate waves of propagating the assembly with actin. Also involved in mechanisms WAVE independent to regulate myosin and actin polymerization during neutrophil chemotaxis (By similarity).[UniProtKB/Swiss-Prot Function]