

Product datasheet for MR211698L4

Ddb1 (NM_015735) Mouse Tagged Lenti ORF Clone

Product data:

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product Type:	Expression Plasmids						
Product Name:	Ddb1 (NM_015735) Mouse Tagged Lenti ORF Clone						
Tag:	mGFP						
Symbol:	Ddb1						
Synonyms:	127kDa; AA408517; p127-Ddb1						
Mammalian Cell Selection:	Puromycin						
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)						
E. coli Selection:	Chloramphenicol (34 ug/mL)						
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR211698).						
Restriction Sites:	Sgfl-Mlul						
Cloning Scheme:							
	Cloning sites used for ORF Shuttling: Sgf 1 ORF Mlu 1 GCG ATC GC ATG // NNN ACG CGT						

						_	Koza Conse		_			
EcoR I	BamH	11	RI	35			S	gf I			OF	RF
CTATAGGGCGGCCGGGAATTCGTC	GACTGGAT	CCGGTAC	CGAG	GAGA	тстб	ccGC	cGCG	ATCG	C C A1	'G	 	
	Mlu I		Notl Xhol			mGFP Tag						
NNŇ	ACG CGT T R	ACG CGO T R	G CCG P	стс L	GAG E	ATG M	AGC S	GGG G	GGC G		 	
GGA CTC AGA GTT TGG G L R V	GTA GGA A	GC										

ACCN: ORF Size: NM_015735 3420 bp



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

* The last codon before the Stop codon of the ORF.

	db1 (NM_015735) Mouse Tagged Lenti ORF Clone – MR211698L4
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Met	 hod: 1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	<u>NM 015735.1</u>
RefSeq Size:	4171 bp
RefSeq ORF:	3423 bp
Locus ID:	13194
UniProt ID:	<u>Q3U1J4</u>
Cytogenetics:	19 6.66 cM

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

GRIGENE Ddb1 (NM_015735) Mouse Tagged Lenti ORF Clone – MR211698L4

Required for DNA repair. Binds to DDB2 to form the UV-damaged DNA-binding protein Gene Summary: complex (the UV-DDB complex). The UV-DDB complex may recognize UV-induced DNA damage and recruit proteins of the nucleotide excision repair pathway (the NER pathway) to initiate DNA repair. The UV-DDB complex preferentially binds to cyclobutane pyrimidine dimers (CPD), 6-4 photoproducts (6-4 PP), apurinic sites and short mismatches. Also appears to function as a component of numerous distinct DCX (DDB1-CUL4-X-box) E3 ubiquitinprotein ligase complexes which mediate the ubiquitination and subsequent proteasomal degradation of target proteins. The functional specificity of the DCX E3 ubiquitin-protein ligase complex is determined by the variable substrate recognition component recruited by DDB1. DCX(DDB2) (also known as DDB1-CUL4-ROC1, CUL4-DDB-ROC1 and CUL4-DDB-RBX1) may ubiguitinate histone H2A, histone H3 and histone H4 at sites of UV-induced DNA damage. The ubiquitination of histones may facilitate their removal from the nucleosome and promote subsequent DNA repair. DCX(DDB2) also ubiquitinates XPC, which may enhance DNA-binding by XPC and promote NER. DCX(DTL) plays a role in PCNA-dependent polyubiquitination of CDT1 and MDM2-dependent ubiquitination of TP53 in response to radiation-induced DNA damage and during DNA replication. DCX(ERCC8) (the CSA complex) plays a role in transcription-coupled repair (TCR). May also play a role in ubiquitination of CDKN1B/p27kip when associated with CUL4 and SKP2 (By similarity). The DDB1-CUL4A-DTL E3 ligase complex regulates the circadian clock function by mediating the ubiquitination and degradation of CRY1 (PubMed:26431207). DDB1-mediated CRY1 degradation promotes FOXO1 protein stability and FOXO1-mediated gluconeogenesis in the liver (PubMed:28790135).[UniProtKB/Swiss-Prot Function]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US