

## Product datasheet for MR211724L3V

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Washc5 (NM 153548) Mouse Tagged ORF Clone Lentiviral Particle

### **Product data:**

**Product Type:** Lentiviral Particles

**Product Name:** Washc5 (NM\_153548) Mouse Tagged ORF Clone Lentiviral Particle

Symbol:

AL022848; C76463; E430025E21Rik; Kiaa0196; mKIAA0196; strumpellin Synonyms:

**Mammalian Cell** 

Selection:

ACCN:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK NM 153548

**ORF Size:** 3477 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(MR211724).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: NM 153548.2

RefSeq Size: 4038 bp RefSeq ORF: 3480 bp Locus ID: 223593 **UniProt ID:** Q8C2E7 Cytogenetics: 15 D1







### **Gene Summary:**

Acts at least in part as component of the WASH core complex whose assembly at the surface of endosomes seems to inhibit WASH nucleation-promoting factor (NPF) activity in recruiting and activating the Arp2/3 complex to induce actin polymerization, and which is involved in regulation of the fission of tubules that serve as transport intermediates during endosome sorting. May be involved in axonal outgrowth. Involved in cellular localization of ADRB2. Involved in cellular trafficking of BLOC-1 complex cargos such as ATP7A and VAMP7 (By similarity). Involved in cytokinesis and following polar body extrusion during oocyte meiotic maturation (PubMed:24998208).[UniProtKB/Swiss-Prot Function]