

Product datasheet for MR211916L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Mapk8ip3 (NM_001163449) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Mapk8ip3 (NM_001163449) Mouse Tagged ORF Clone Lentiviral Particle

Symbol: Mapk8ip3

Synonyms: BB120594; D17Wsu15e; JIP-3; Jip3; JSAP1; JSAP1a; JSAP1b; JSAP1c; JSAP1d; mKIAA1066; Syd2

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK

ACCN: NM_001163449

ORF Size: 3942 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(MR211916).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 001163449.1</u>, <u>NP 001156921.1</u>

RefSeq Size: 5510 bp
RefSeq ORF: 3945 bp
Locus ID: 30957
UniProt ID: Q9ESN9

Cytogenetics: 17 12.53 cM





Gene Summary:

The JNK-interacting protein (JIP) group of scaffold proteins selectively mediates JNK signaling by aggregating specific components of the MAPK cascade to form a functional JNK signaling module. May function as a regulator of vesicle transport, through interactions with the JNKsignaling components and motor proteins (PubMed:10523642, PubMed:10629060). Promotes neuronal axon elongation in a kinesin- and JNK-dependent manner (PubMed:23576431, PubMed:25944905, PubMed:28259553). Activates cofilin at axon tips via local activation of INK, thereby regulating filopodial dynamics and enhancing axon elongation (PubMed:23576431, PubMed:25944905, PubMed:28259553). Its binding to kinesin heavy chains (KHC), promotes kinesin-1 motility along microtubules and is essential for axon elongation and regeneration (PubMed:23576431, PubMed:25944905, PubMed:28259553). Regulates cortical neuronal migration by mediating NTRK2/TRKB anterograde axonal transport during brain development (PubMed:23576431, PubMed:25944905, PubMed:28259553). Acts as an adapter that bridges the interaction between NTRK2/TRKB and KLC1 and drives NTRK2/TRKB axonal but not dendritic anterograde transport, which is essential for subsequent BDNF-triggered signaling and filopodia formation (PubMed:23576431, PubMed:25944905, PubMed:28259553).[UniProtKB/Swiss-Prot Function]