

## Product datasheet for MR211946L4V

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Usp19 (NM\_027804) Mouse Tagged ORF Clone Lentiviral Particle

### **Product data:**

Product Type: Lentiviral Particles

**Product Name:** Usp19 (NM\_027804) Mouse Tagged ORF Clone Lentiviral Particle

Symbol: Usp19

**Synonyms:** 8430421107Rik; Al047774; Zmynd9

**Mammalian Cell** 

Selection:

Puromycin

**Vector:** pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

**ACCN:** NM\_027804 **ORF Size:** 4080 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(MR211946).

Sequence:
OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 027804.4, NP 082080.3

 RefSeq Size:
 4727 bp

 RefSeq ORF:
 4083 bp

 Locus ID:
 71472

 UniProt ID:
 Q3UJD6

 Cytogenetics:
 9 F2







### **Gene Summary:**

Deubiquitinating enzyme that regulates the degradation of various proteins. Deubiquitinates and prevents proteasomal degradation of RNF123 which in turn stimulates CDKN1B ubiquitin-dependent degradation thereby playing a role in cell proliferation. Involved in decreased protein synthesis in atrophying skeletal muscle. Modulates transcription of major myofibrillar proteins. Also involved in turnover of endoplasmic-reticulum-associated degradation (ERAD) substrates (By similarity). Regulates the stability of BIRC2/c-IAP1 and BIRC3/c-IAP2 by preventing thier ubiquitination. Required for cells to mount an appropriate response to hypoxia and rescues HIF1A from degradation in a non-catalytic manner. Exhibits a preference towards 'Lys-63'-linked ubiquitin chains (By similarity). Plays an important role in 17 beta-estradiol (E2)-inhibited myogenesis. Decreases the levels of ubiquitinated proteins during skeletal muscle formation and acts to repress myogenesis. [UniProtKB/Swiss-Prot Function]