

## Product datasheet for MR211949L4V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Med23 (NM\_027347) Mouse Tagged ORF Clone Lentiviral Particle

**Product data:** 

Product Type: Lentiviral Particles

**Product Name:** Med23 (NM\_027347) Mouse Tagged ORF Clone Lentiviral Particle

Symbol: Med23

**Synonyms:** 130kDa; 3000002A17Rik; Crsp3; ESTM7; mKIAA1216; Sur2; X83317

Mammalian Cell

Selection:

Puromycin

**Vector:** pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

**ACCN:** NM\_027347 **ORF Size:** 4110 bp

**ORF Nucleotide** 

•

Sequence:

The ORF insert of this clone is exactly the same as(MR211949).

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This

clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 027347.3, NP 081623.3

RefSeq Size: 4928 bp
RefSeq ORF: 4113 bp
Locus ID: 70208
UniProt ID: Q80YQ2

Cytogenetics: 10







## **Gene Summary:**

Component of the Mediator complex, a coactivator involved in the regulated transcription of nearly all RNA polymerase II-dependent genes. Mediator functions as a bridge to convey information from gene-specific regulatory proteins to the basal RNA polymerase II transcription machinery. Mediator is recruited to promoters by direct interactions with regulatory proteins and serves as a scaffold for the assembly of a functional pre-initiation complex with RNA polymerase II and the general transcription factors (By similarity). Also required for transcriptional activation subsequent to the assembly of the pre-initiation complex. Required for transcriptional activation by adenovirus E1A protein. Required for ELK1-dependent transcriptional activation in response to activated Ras signaling. [UniProtKB/Swiss-Prot Function]