

Product datasheet for MR211961L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Prx (BC068135) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Prx (BC068135) Mouse Tagged ORF Clone Lentiviral Particle

Symbol:

L-Periaxin Synonyms: **Mammalian Cell**

Selection:

Puromycin

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

mGFP Tag:

ACCN: BC068135 **ORF Size:** 4173 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(MR211961).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: BC068135.1 RefSeq Size: 4502 bp RefSeq ORF: 4175 bp Locus ID: 19153

Cytogenetics: 7 15.91 cM







Gene Summary:

Scaffolding protein that functions as part of a dystroglycan complex in Schwann cells, and as part of EZR and AHNAK-containing complexes in eye lens fiber cells (PubMed:11430802, PubMed:21745462, PubMed:22764250). Required for the maintenance of the peripheral myelin sheath that is essential for normal transmission of nerve impulses and normal perception of sensory stimuli (PubMed:10839370). Required for normal transport of MBP mRNA from the perinuclear to the paranodal regions (PubMed:15356632). Required for normal remyelination after nerve injury (PubMed:10839370). Required for normal elongation of Schwann cells and normal length of the internodes between the nodes of Ranvier. The demyelinated nodes of Ranvier permit saltatory transmission of nerve impulses; shorter internodes cause slower transmission of nerve impulses (PubMed:15356632, PubMed:23022068). Required for the formation of appositions between the abaxonal surface of the myelin sheath and the Schwann cell plasma membrane; the Schwann cell cytoplasm is restricted to regions between these appositions (PubMed:15356632, PubMed:23022068). Required for the formation of Cajal bands and of Schmidt-Lanterman incisures that correspond to short, cytoplasm-filled regions on myelinated nerves (PubMed:23022068, PubMed:22764250). Recruits DRP2 to the Schwann cell plasma membrane (PubMed:11430802, PubMed:23022068, PubMed:22764250). Required for normal protein composition of the eye lens fiber cell plasma membrane and normal eye lens fiber cell morphology (PubMed:21745462).[UniProtKB/Swiss-Prot Function]