

Product datasheet for MR212025L3

Eif2ak4 (NM_001177806) Mouse Tagged Lenti ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Eif2ak4 (NM_001177806) Mouse Tagged Lenti ORF Clone
Tag:	Myc-DDK
Symbol:	Eif2ak4
Synonyms:	2610011M03; GCN2; MGCN2
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
E. coli Selection:	Chloramphenicol (34 ug/mL)
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR212025).
Restriction Sites:	SgfI-MluI
Cloning Scheme:	


Cloning sites used for ORF Shuttling:



* The last codon before the Stop codon of the ORF.



[View online »](#)

Plasmid Map:


ACCN: NM_001177806

ORF Size: 4611 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_001177806.1](#), [NP_001171277.1](#)

RefSeq Size: 4916 bp

RefSeq ORF: 4611 bp

Locus ID: 27103

UniProt ID: [Q9QZ05](#)

Cytogenetics: 2 E5

Gene Summary: Metabolic-stress sensing protein kinase that phosphorylates the alpha subunit of eukaryotic translation initiation factor 2 (eIF-2-alpha/EIF2S1) on 'Ser-52' in response to low amino acid availability (PubMed:10504407, PubMed:10655230, PubMed:12176355, PubMed:12215525, PubMed:15213227, PubMed:16054071, PubMed:16176978, PubMed:16121183, PubMed:15774759, PubMed:16601681, PubMed:26102367). Plays a role as an activator of the integrated stress response (ISR) required for adaptation to amino acid starvation. Converts phosphorylated eIF-2-alpha/EIF2S1 either to a competitive inhibitor of the translation initiation factor eIF-2B, leading to a global protein synthesis repression, and thus to a reduced overall utilization of amino acids, or to a translational initiation activation of specific mRNAs, such as the transcriptional activator ATF4, and hence allowing ATF4-mediated reprogramming of amino acid biosynthetic gene expression to alleviate nutrient depletion (PubMed:10655230, PubMed:11106749, PubMed:12176355, PubMed:15213227, PubMed:16176978, PubMed:26102367). Binds uncharged tRNAs (By similarity). Involved in cell cycle arrest by promoting cyclin D1 mRNA translation repression after the unfolded protein response pathway (UPR) activation or cell cycle inhibitor CDKN1A/p21 mRNA translation activation in response to amino acid deprivation (PubMed:16176978, PubMed:26102367). Plays a role in the consolidation of synaptic plasticity, learning as well as formation of long-term memory (PubMed:16121183). Plays a role in neurite outgrowth inhibition (PubMed:23447528). Plays a role in feeding behavior to maintain amino acid homeostasis; contributes to the innate aversion toward diets of imbalanced amino acid composition (PubMed:16054071, PubMed:15774759). Plays a proapoptotic role in response to glucose deprivation (PubMed:20660158). Promotes global cellular protein synthesis repression in response to UV irradiation independently of the stress-activated protein kinase/c-Jun N-terminal kinase (SAPK/JNK) and p38 MAPK signaling pathways (PubMed:12176355).[UniProtKB/Swiss-Prot Function]