

Product datasheet for **MR212025L4V**

Eif2ak4 (NM_001177806) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	Eif2ak4 (NM_001177806) Mouse Tagged ORF Clone Lentiviral Particle
Symbol:	Eif2ak4
Synonyms:	2610011M03; GCN2; MGCN2
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
Tag:	mGFP
ACCN:	NM_001177806
ORF Size:	4611 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR212025).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	NM_001177806.1 , NP_001171277.1
RefSeq Size:	4916 bp
RefSeq ORF:	4611 bp
Locus ID:	27103
UniProt ID:	Q9QZ05
Cytogenetics:	2 E5



[View online »](#)

Gene Summary:

Metabolic-stress sensing protein kinase that phosphorylates the alpha subunit of eukaryotic translation initiation factor 2 (eIF-2-alpha/EIF2S1) on 'Ser-52' in response to low amino acid availability (PubMed:10504407, PubMed:10655230, PubMed:12176355, PubMed:12215525, PubMed:15213227, PubMed:16054071, PubMed:16176978, PubMed:16121183, PubMed:15774759, PubMed:16601681, PubMed:26102367). Plays a role as an activator of the integrated stress response (ISR) required for adaptation to amino acid starvation. Converts phosphorylated eIF-2-alpha/EIF2S1 either to a competitive inhibitor of the translation initiation factor eIF-2B, leading to a global protein synthesis repression, and thus to a reduced overall utilization of amino acids, or to a translational initiation activation of specific mRNAs, such as the transcriptional activator ATF4, and hence allowing ATF4-mediated reprogramming of amino acid biosynthetic gene expression to alleviate nutrient depletion (PubMed:10655230, PubMed:11106749, PubMed:12176355, PubMed:15213227, PubMed:16176978, PubMed:26102367). Binds uncharged tRNAs (By similarity). Involved in cell cycle arrest by promoting cyclin D1 mRNA translation repression after the unfolded protein response pathway (UPR) activation or cell cycle inhibitor CDKN1A/p21 mRNA translation activation in response to amino acid deprivation (PubMed:16176978, PubMed:26102367). Plays a role in the consolidation of synaptic plasticity, learning as well as formation of long-term memory (PubMed:16121183). Plays a role in neurite outgrowth inhibition (PubMed:23447528). Plays a role in feeding behavior to maintain amino acid homeostasis; contributes to the innate aversion toward diets of imbalanced amino acid composition (PubMed:16054071, PubMed:15774759). Plays a proapoptotic role in response to glucose deprivation (PubMed:20660158). Promotes global cellular protein synthesis repression in response to UV irradiation independently of the stress-activated protein kinase/c-Jun N-terminal kinase (SAPK/JNK) and p38 MAPK signaling pathways (PubMed:12176355). [UniProtKB/Swiss-Prot Function]