

Product datasheet for MR212048L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Eif4g1 (NM_145941) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Eif4g1 (NM_145941) Mouse Tagged ORF Clone Lentiviral Particle

Symbol: Eif4g⁻

Synonyms: E030015G23Rik; elF-4-gamma 1; elF-4G 1; elF-4G1; elF4Gl

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_145941 **ORF Size:** 4800 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(MR212048).

Sequence:
OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 145941.2, NP 666053.2

RefSeq Size: 5481 bp
RefSeq ORF: 4803 bp
Locus ID: 208643
UniProt ID: Q6NZJ6
Cytogenetics: 16 B1







Gene Summary:

This gene encodes a member of the eukaryotic translation initiation factors (eIF) that play important roles in translation initiation by mediating recruitment of additional initiation factors and providing a scaffold for ribosome/mRNA-bridging. Along with eIF4A and eIF4E, the encoded protein forms the eIF4F complex that bridges the 5' UTR with the polyadenylated 3' UTR resulting in mRNA circularization, enhanced translation initiation and mRNA stability. Through its association with eIF3, the encoded protein mediates recruitment of the 43S preinitiation complex to mRNA. Alternative splicing of this gene results in multiple transcript variants. Pseudogenes for this gene have been identified on chromosomes 2 and 13. [provided by RefSeq, Jan 2015]