

## Product datasheet for MR212701L3

### Hpf1 (NM\_028299) Mouse Tagged Lenti ORF Clone

#### Product data:

Product Type:	Expression Plasmids
Product Name:	Hpf1 (NM_028299) Mouse Tagged Lenti ORF Clone
Tag:	Myc-DDK
Symbol:	Hpf1
Synonyms:	2700029M09Rik; C4orf27; C230006B22Rik
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
E. coli Selection:	Chloramphenicol (34 ug/mL)
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR212701).
Restriction Sites:	SgfI-MluI
Cloning Scheme:	

Cloning sites used for ORF Shuttling:



\* The last codon before the Stop codon of the ORF.


ACCN:	NM_028299
ORF Size:	1038 bp



[View online »](#)

<b>OTI Disclaimer:</b>	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a>
<b>OTI Annotation:</b>	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
<b>Components:</b>	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
<b>Reconstitution Method:</b>	<ol style="list-style-type: none"><li>1. Centrifuge at 5,000xg for 5min.</li><li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li><li>3. Close the tube and incubate for 10 minutes at room temperature.</li><li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li><li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li></ol>
<b>RefSeq:</b>	<a href="#">NM_028299.1</a>
<b>RefSeq Size:</b>	1210 bp
<b>RefSeq ORF:</b>	1041 bp
<b>Locus ID:</b>	72612
<b>UniProt ID:</b>	<a href="#">Q8CFE2</a>
<b>Cytogenetics:</b>	8 B3.1
<b>Gene Summary:</b>	Acts as a cofactor for serine ADP-ribosylation by conferring serine specificity on PARP1 and PARP2: interacts with PARP1 and PARP2 and is able to change amino acid specificity toward serine. Promotes histone serine ADP-ribosylation in response to DNA damage, limiting DNA damage-induced PARP1 hyper-automodification, and ensuring genome stability. Serine ADP-ribosylation of proteins constitutes the primary form of ADP-ribosylation of proteins in response to DNA damage. HPF1 also promotes tyrosine ADP-ribosylation, probably by conferring tyrosine specificity on PARP1.[UniProtKB/Swiss-Prot Function]

Product images:



Circular map for MR212701L3