

Product datasheet for MR214240L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Ythdc1 (NM_177680) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Ythdc1 (NM 177680) Mouse Tagged ORF Clone Lentiviral Particle

Symbol: Ythdc1

Synonyms: A730098D12Rik; C80342; mKIAA1966

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

 Tag:
 Myc-DDK

 ACCN:
 NM_177680

ORF Size: 2208 bp

ORF Nucleotide

Sequence:

The ORF insert of this clone is exactly the same as(MR214240).

OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 177680.3

 RefSeq Size:
 3034 bp

 RefSeq ORF:
 2211 bp

 Locus ID:
 231386

 UniProt ID:
 E9Q5K9

 Cytogenetics:
 5 E1





Gene Summary:

Regulator of alternative splicing that specifically recognizes and binds N6-methyladenosine (m6A)-containing RNAs (By similarity). M6A is a modification present at internal sites of mRNAs and some non-coding RNAs and plays a role in the efficiency of mRNA splicing, processing and stability (By similarity). Acts as a key regulator of exon-inclusion or exonskipping during alternative splicing via interaction with mRNA splicing factors SRSF3 and SRSF10 (By similarity). Specifically binds m6A-containing mRNAs and promotes recruitment of SRSF3 to its mRNA-binding elements adjacent to m6A sites, leading to exon-inclusion during alternative splicing (By similarity). In contrast, interaction with SRSF3 prevents interaction with SRSF10, a splicing factor that promotes exon skipping: this prevents SRSF10 from binding to its mRNA-binding sites close to m6A-containing regions, leading to inhibit exon skipping during alternative splicing (By similarity). May also regulate alternative splice site selection (By similarity). Also involved in nuclear export of m6A-containing mRNAs via interaction with SRSF3: interaction with SRSF3 facilitates m6A-containing mRNA-binding to both SRSF3 and NXF1, promoting mRNA nuclear export (By similarity). Also recognizes and binds m6A on other RNA molecules (By similarity). Involved in random X inactivation mediated by Xist RNA: recognizes and binds m6A-containing Xist and promotes transcription repression activity of Xist (By similarity). Involved in S-adenosyl-L-methionine homeostasis by regulating expression of MAT2A transcripts, probably by binding m6A-containing MAT2A mRNAs (PubMed:29262316).[UniProtKB/Swiss-Prot Function]