

Product datasheet for MR214444L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Thoc2 (NM_001033422) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Thoc2 (NM 001033422) Mouse Tagged ORF Clone Lentiviral Particle

Symbol: Thoc2

Synonyms: 6330441O12Rik; D130005M13Rik; Gm1139; Gm1793; Tho2

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_001033422

ORF Size: 4782 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(MR214444).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: NM 001033422.1, NP 001028594.1

RefSeq Size: 7631 bp
RefSeq ORF: 4785 bp
Locus ID: 331401
UniProt ID: B1AZI6
Cytogenetics: X A4







Gene Summary:

Required for efficient export of polyadenylated RNA and spliced mRNA. Acts as component of the THO subcomplex of the TREX complex which is thought to couple mRNA transcription, processing and nuclear export, and which specifically associates with spliced mRNA and not with unspliced pre-mRNA. TREX is recruited to spliced mRNAs by a transcription-independent mechanism, binds to mRNA upstream of the exon-junction complex (EJC) and is recruited in a splicing- and cap-dependent manner to a region near the 5' end of the mRNA where it functions in mRNA export to the cytoplasm via the TAP/NFX1 pathway. Plays a role for proper neuronal development.[UniProtKB/Swiss-Prot Function]