

## Product datasheet for MR215161L3V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Abcc4 (NM\_001163675) Mouse Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

**Product Name:** Abcc4 (NM\_001163675) Mouse Tagged ORF Clone Lentiviral Particle

Symbol: Abcc4

Synonyms: ABCC4-N1; D630049P08Rik; MOATB; MRP4

Mammalian Cell

Selection:

Puromycin

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK

**ACCN:** NM\_001163675

ORF Size: 3975 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(MR215161).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This

clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 001163675.1

 RefSeq Size:
 5711 bp

 RefSeq ORF:
 3849 bp

 Locus ID:
 239273

 UniProt ID:
 Q3TZN9

Cytogenetics: 14 E4





## **Gene Summary:**

ATP-dependent transporter of the ATP-binding cassette (ABC) family that actively extrudes physiological compounds and xenobiotics from cells. Transports a range of endogenous molecules that have a key role in cellular communication and signaling, including cyclic nucleotides such as cyclic AMP (cAMP) and cyclic GMP (cGMP), bile acids, steroid conjugates, urate, and prostaglandins. Mediates also the ATP-dependent efflux of glutathione conjugates such as leukotriene C4 (LTC4) and leukotriene B4 (LTB4). The presence of GSH is necessary for the ATP-dependent transport of LTB4, whereas GSH is not required for the transport of LTC4. Mediates the cotransport of bile acids with reduced glutathione (GSH). Transports a wide range of drugs and their metabolites, including anticancer, antiviral and antibiotics molecules (Probable). Confers resistance to anticancer agents (Probable). [UniProtKB/Swiss-Prot Function]