

# Product datasheet for MR215388

### H2bw2 (NM\_027067) Mouse Tagged ORF Clone

### **Product data:**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product Type:	Expression Plasmids
Product Name:	H2bw2 (NM_027067) Mouse Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	H2bw2
Synonyms:	1700014N06Rik; H2bfm; H2bfw; H2BL; H2bl2
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
ORF Nucleotide Sequence:	>MR215388 ORF sequence Red=Cloning site Blue=ORF Green=Tags(s)
	TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC GCC <mark>GCGATCGC</mark> C

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT ACAAGGATGACGACGATAAGGTTTAA



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

Duetein Comment	NMP215200 protoin sequence
Protein Sequence:	>MR215388 protein sequence Red=Cloning site Green=Tags(s)
	MASTTAMDVLEELSSDSSEKQVQPRKPEKAKREKDKPKKGGPEKKAKKEKQEKAKPEKKPKKKPEKEKPE GEKLEKKPKKDKREKAKPKKKPEQENREQETPEQEKPEVQRRRSLHQSIREDERRARLIRRRKNSFAIYF PKVLKNIHVGLSLSQRSVNILDSFVKDMFERIASEASFLARQARNSTINSREIQTAIRLLLPGELCRRAV AEGTMAMVRYISNK
	TRTRPLEQKLISEEDLAANDILDYKDDDDKV
<b>Restriction Sites:</b>	Sgfl-Mlul
Cloning Scheme:	Cloning sites used for ORF Shuttling:
	Sgfi ORF Miui GCGATOGC C ATG// NIN ACG CGT
	Kozac         Consensus         EcoRI       BamHI Kpn I       RBS       SgfI         CTATASGGCGCCCGGGAATTCCCGTACCGAGGAGGATCTCGCCCGCGGCGATCGC C       RTC
	ORF <u>Miu I Not I Xho I</u> Myc.Tag ACG CGT ACG CGG CCG CTC GAG CAG AAA CTC ATC TCA GAA GAG T R T R P L E Q K L I S E E
	EcoR V     Flag.Tag     Pme /     Fse /       GAT CTG GCA GCA AAT GAT ATC CTG GAT TAC AAG GAT GAC GAC GAT AAG GTT TAA ACGGCCGGCC     D     L     A     N     D     I     L     D     Y     K     D     D     D     K     V     stop
	* The last codon before the Stop codon of the ORF
ACCN:	NM_027067
ORF Size:	675 bp
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).


This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US 

### 

#### H2bw2 (NM\_027067) Mouse Tagged ORF Clone - MR215388

Reconstitution Method:	<ol> <li>Centrifuge at 5,000xg for 5min.</li> <li>Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li> <li>Close the tube and incubate for 10 minutes at room temperature.</li> <li>Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li> <li>Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li> </ol>
RefSeq:	<u>NM 027067.2, NP 081343.1</u>
RefSeq Size:	1049 bp
RefSeq ORF:	675 bp
Locus ID:	69389
Cytogenetics:	X F1
MW:	26.1 kDa
Gene Summary:	Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chromatin into higher order structures. This gene encodes a replication-independent histone that is a member of the H2B histone family. [provided by RefSeq, Nov 2015]

## **Product images:**



Circular map for MR215388

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US