

Product datasheet for MR215666L4

Loxl2 (NM_033325) Mouse Tagged Lenti ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: Loxl2 (NM_033325) Mouse Tagged Lenti ORF Clone

Tag: mGFP Symbol: Loxl2

Synonyms: 1110004B06Rik; 4930526G11Rik; 9430067E15Rik

Mammalian Cell Puromycin

Selection:

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

E. coli Selection: Chloramphenicol (34 ug/mL)


ORF Nucleotide The ORF insert of this clone is exactly the same as(MR215666).

Sequence:

Restriction Sites: Sgfl-Mlul

Cloning Scheme:





^{*} The last codon before the Stop codon of the ORF.

ACCN: NM_033325

ORF Size: 2328 bp



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



LoxI2 (NM_033325) Mouse Tagged Lenti ORF Clone - MR215666L4

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 033325.2</u>, <u>NP 201582.2</u>

P58022

 RefSeq Size:
 5186 bp

 RefSeq ORF:
 2331 bp

 Locus ID:
 94352

Cytogenetics: 14 D2


UniProt ID:



Gene Summary:

Mediates the post-translational oxidative deamination of lysine residues on target proteins leading to the formation of deaminated lysine (allysine) (By similarity). Acts as a transcription corepressor and specifically mediates deamination of trimethylated 'Lys-4' of histone H3 (H3K4me3), a specific tag for epigenetic transcriptional activation (By similarity). Shows no activity against histone H3 when it is trimethylated on 'Lys-9' (H3K9me3) or 'Lys-27' (H3K27me3) or when 'Lys-4' is monomethylated (H3K4me1) or dimethylated (H3K4me2) (By similarity). Also mediates deamination of methylated TAF10, a member of the transcription factor IID (TFIID) complex, which induces release of TAF10 from promoters, leading to inhibition of TFIID-dependent transcription (By similarity). LOXL2-mediated deamination of TAF10 results in transcriptional repression of genes required for embryonic stem cell pluripotency including POU5F1/OCT4, NANOG, KLF4 and SOX2 (PubMed:25959397). Involved in epithelial to mesenchymal transition (EMT) via interaction with SNAI1 and participates in repression of E-cadherin, probably by mediating deamination of histone H3 (By similarity). During EMT, involved with SNAI1 in negatively regulating pericentromeric heterochromatin transcription (By similarity). SNAI1 recruits LOXL2 to pericentromeric regions to oxidize histone H3 and repress transcription which leads to release of heterochromatin component CBX5/HP1A, enabling chromatin reorganization and acquisition of mesenchymal traits (By similarity). Interacts with the endoplasmic reticulum protein HSPA5 which activates the IRE1-XBP1 pathway of the unfolded protein response, leading to expression of several transcription factors involved in EMT and subsequent EMT induction (By similarity). When secreted into the extracellular matrix, promotes cross-linking of extracellular matrix proteins by mediating oxidative deamination of peptidyl lysine residues in precursors to fibrous collagen and elastin (By similarity). Acts as a regulator of sprouting angiogenesis, probably via collagen IV scaffolding (By similarity). Acts as a regulator of chondrocyte differentiation, probably by regulating expression of factors that control chondrocyte differentiation (PubMed:21071451).[UniProtKB/Swiss-Prot Function]

Product images:



Circular map for MR215666L4