

Product datasheet for MR216075L4

Calcoco2 (NM_029755) Mouse Tagged Lenti ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Calcoco2 (NM_029755) Mouse Tagged Lenti ORF Clone
Tag:	mGFP
Symbol:	Calcoco2
Synonyms:	2410154J16Rik; C77254; Ndp52; Ndp52l1
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
E. coli Selection:	Chloramphenicol (34 ug/mL)
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR216075).
Restriction Sites:	SgfI-MluI
Cloning Scheme:	


Cloning sites used for ORF Shuttling:



* The last codon before the Stop codon of the ORF.



[View online »](#)

Plasmid Map:


ACCN: NM_029755

ORF Size: 930 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_029755.3](#), [NP_084031.2](#)

RefSeq Size: 1556 bp

RefSeq ORF: 933 bp

Locus ID: 76815

Cytogenetics: 11 59.4 cM

Gene Summary: Xenophagy-specific receptor required for autophagy-mediated intracellular bacteria degradation (By similarity). Acts as an effector protein of galectin-sensed membrane damage that restricts the proliferation of infecting pathogens upon entry into the cytosol by targeting LGALS8-associated bacteria for autophagy (By similarity). Initially orchestrates bacteria targeting to autophagosomes and subsequently ensures pathogen degradation by regulating pathogen-containing autophagosome maturation (By similarity). Bacteria targeting to autophagosomes relies on its interaction with MAP1LC3A, MAP1LC3B and/or GABARAPL2, whereas regulation of pathogen-containing autophagosome maturation requires the interaction with MAP3LC3C (By similarity). May play a role in ruffle formation and actin cytoskeleton organization and seems to negatively regulate constitutive secretion (By similarity).[UniProtKB/Swiss-Prot Function]