

## Product datasheet for MR216083L3V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Osbp2 (NM\_152818) Mouse Tagged ORF Clone Lentiviral Particle

**Product data:** 

Product Type: Lentiviral Particles

**Product Name:** Osbp2 (NM 152818) Mouse Tagged ORF Clone Lentiviral Particle

Symbol: Osbp2

**Synonyms:** 1700095P05Rik; C630001G20Rik; Gm244; ORP-4; OSBPL1

**Mammalian Cell** 

Selection:

Puromycin

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK
ACCN: NM 152818

ORF Size: 2724 bp

**ORF Nucleotide** 

ucleotide The O

Sequence:
OTI Disclaimer:

The ORF insert of this clone is exactly the same as(MR216083).

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 152818.2</u>

 RefSeq Size:
 4138 bp

 RefSeq ORF:
 2727 bp

 Locus ID:
 74309

 UniProt ID:
 Q5QNQ6

Cytogenetics: 11 A1







## **Gene Summary:**

The protein encoded by this gene belongs to the oxysterol-binding protein-related family of proteins, which are defined by a C-terminal sterol domain with a highly conserved EQVSHHPP motif. Oxysterols are oxygenated derivatives of cholesterol that are involved in mechanisms that include apoptosis, cholesterol homeostasis, lipid trafficking and cell differentiation. This protein is selectively expressed at high levels in the brain and testis. Within the testis, the mRNA is localized to postmeiotic germ cells, including spermatids and spermatozoa, but is not detectable in somatic cells. Mice homozygous mutant for a targeted deletion in this gene do not exhibit overt developmental phenotypes but are male sterile. Females display normal fertility. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Oct 2014]