

## Product datasheet for **MR216271L3V**

### Rnf20 (NM\_182999) Mouse Tagged ORF Clone Lentiviral Particle

#### Product data:

Product Type:	Lentiviral Particles
Product Name:	Rnf20 (NM_182999) Mouse Tagged ORF Clone Lentiviral Particle
Symbol:	Rnf20
Synonyms:	4833430L21Rik; AW540162; C79397; mKIAA4116
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
Tag:	Myc-DDK
ACCN:	NM_182999
ORF Size:	2919 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR216271).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<a href="#">NM_182999.2</a> , <a href="#">NP_892044.1</a>
RefSeq Size:	4159 bp
RefSeq ORF:	2922 bp
Locus ID:	109331
UniProt ID:	<a href="#">Q5DTM8</a>
Cytogenetics:	4 B1



[View online »](#)

**Gene Summary:**

Component of the RNF20/40 E3 ubiquitin-protein ligase complex that mediates monoubiquitination of 'Lys-120' of histone H2B (H2BK120ub1). H2BK120ub1 gives a specific tag for epigenetic transcriptional activation and is also prerequisite for histone H3 'Lys-4' and 'Lys-79' methylation (H3K4me and H3K79me, respectively). It thereby plays a central role in histone code and gene regulation. The RNF20/40 complex forms a H2B ubiquitin ligase complex in cooperation with the E2 enzyme UBE2A or UBE2B; reports about the cooperation with UBE2E1/UBCH are contradictory. Required for transcriptional activation of Hox genes. Recruited to the MDM2 promoter, probably by being recruited by p53/TP53, and thereby acts as a transcriptional coactivator. Mediates the polyubiquitination of PA2G4 leading to its proteasome-mediated degradation.[UniProtKB/Swiss-Prot Function]