

Product datasheet for MR216659L4

Ppie (NM_019489) Mouse Tagged Lenti ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Ppie (NM_019489) Mouse Tagged Lenti ORF Clone
Tag:	mGFP
Symbol:	Ppie
Synonyms:	2010010D16Rik; Cyp33
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
E. coli Selection:	Chloramphenicol (34 ug/mL)
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR216659).
Restriction Sites:	SgfI-MluI
Cloning Scheme:	

Cloning sites used for ORF Shuttling:



* The last codon before the Stop codon of the ORF.


ACCN:	NM_019489
ORF Size:	903 bp



[View online »](#)

OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_019489.5
RefSeq Size:	1168 bp
RefSeq ORF:	906 bp
Locus ID:	56031
UniProt ID:	Q9QZH3
Cytogenetics:	4 D2.2
Gene Summary:	Involved in pre-mRNA splicing as component of the spliceosome. Combines RNA-binding and PPlase activities. Binds mRNA and has a preference for single-stranded RNA molecules with poly-A and poly-U stretches, suggesting it binds to the poly(A)-region in the 3' UTR of mRNA molecules. Catalyzes the cis-trans isomerization of proline imidic peptide bonds in proteins. Inhibits KMT2A activity; this requires proline isomerase activity.[UniProtKB/Swiss-Prot Function]

Product images:



Circular map for MR216659L4