

Product datasheet for MR216803

Rab43 (NM 001039394) Mouse Tagged ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: Rab43 (NM_001039394) Mouse Tagged ORF Clone

Tag: Myc-DDK Symbol: Rab43

Synonyms: 1810048P08Rik; 2500004H21Rik; AW490415

Mammalian Cell Neomycin

Selection:

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

ORF Nucleotide >MR216803 representing NM_001039394
Sequence: Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

ATGGCGGGCCCTGGCCCGGAGATCAGGACGAACACTACGATTTCCTGTTCAAGCTGGTGTTAGTGGGCG
ACGCGAGCGTGGGCAAGACGTGCGTGCTGCAGCGCTTCAAGACCGGCGCCTTCTCGGCGCGTCAGGGCAG
CACCATCGGTGTCGACTTCACCATGAAGACGCTGGAGATCCAGGGCAAGCGGGTCAAGTTACAGATTTGG
GACACAGCTGGCCAGGAGCGGTTCCGGACCATCACCCAGAGCTACTACCGAAGTGCCAATGGGGCTATCC
TTGCGTACGACATCAGCAAGAGGAGCACCTTCTTGTCAGTGCCTCACTGGATCGAGGATGTGAGGAAGTA
CGCGGGTTCCAACATCGTGCAGCTGCTGATTGGGAACAAGTCAGACCTTGCCGATTTCCGGAGGTCCCA
TTGGCGGAGGCACAGAGCCTGGCTGAGCACTATGACATCCTCTGTGCCATCGAGACCTCCGCAAAGGACT
CCAGCAATGTGGAGAGGCCTTCACCAGAGTGGCCACGGAGGTCCCATGTT
CAGTGAGAAGAACACAGACCACATCCAGCTGGACAGCAAGGACATTGCGGAGAGCTGGGGCTTTGGGTGC

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT

ACAAGGATGACGACGATAAGGTTTAA

Protein Sequence: >MR216803 representing NM_001039394

Red=Cloning site Green=Tags(s)

MAGPGPGDQDEHYDFLFKLVLVGDASVGKTCVVQRFKTGAFSARQGSTIGVDFTMKTLEIQGKRVKLQIW DTAGQERFRTITQSYYRSANGAILAYDISKRSTFLSVPHWIEDVRKYAGSNIVQLLIGNKSDLADFREVP LAEAQSLAEHYDILCAIETSAKDSSNVEEAFTRVATELIMRHGGPMFSEKNTDHIQLDSKDIAESWGCGC

TRTRPLEQKLISEEDLAANDILDYKDDDDKV



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Chromatograms: https://cdn.origene.com/chromatograms/mm9060 d10.zip

Restriction Sites: Sgfl-Mlul

Cloning Scheme:





^{*} The last codon before the Stop codon of the ORF

ACCN: NM_001039394

ORF Size: 630 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 001039394.1, NP 001034483.1</u>

RefSeq Size: 4605 bp RefSeq ORF: 633 bp Locus ID: 69834



UniProt ID: Q8CG50

Cytogenetics: 6 D1

MW: 23.3 kDa


Gene Summary: The small GTPases Rab are key regulators of intracellular membrane trafficking, from the

formation of transport vesicles to their fusion with membranes. Rabs cycle between an inactive GDP-bound form and an active GTP-bound form that is able to recruit to membranes different set of downstream effectors directly responsible for vesicle formation, movement, tethering and fusion. The low intrinsic GTPase activity of RAB43 is activated by USP6NL. Involved in retrograde transport from the endocytic pathway to the Golgi apparatus. Involved in the transport of Shiga toxin from early and recycling endosomes to the trans-Golgi

in the transport of Shiga toxin from early and recycling endosomes to the trans-Golgi network. Required for the structural integrity of the Golgi complex. Plays a role in the maturation of phagosomes that engulf pathogens, such as S.aureus and Mycobacterium.

[UniProtKB/Swiss-Prot Function]

Product images:



Circular map for MR216803