

Product datasheet for MR216854L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Sarm1 (NM_001168521) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Sarm1 (NM_001168521) Mouse Tagged ORF Clone Lentiviral Particle

Symbol: Sarm²

Synonyms: A830091I15Rik; C78606; MyD885; Sarm

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK

ACCN: NM_001168521

ORF Size: 2292 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(MR216854).

Sequence:
OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 001168521.1</u>, <u>NP 001161993.1</u>

 RefSeq Size:
 5164 bp

 RefSeq ORF:
 2295 bp

 Locus ID:
 237868

 UniProt ID:
 Q6PDS3

Cytogenetics: 11 46.74 cM







Gene Summary:

Negative regulator of MYD88- and TRIF-dependent toll-like receptor signaling pathway which plays a pivotal role in activating axonal degeneration following injury. Promotes Wallerian degeneration an injury-induced axonal death pathway which involves degeneration of an axon distal to the injury site. Can activate neuronal death in response to stress. Regulates dendritic arborization through the MAPK4-JNK pathway. Involved in innate immune response. Inhibits both TICAM1/TRIF- and MYD88-dependent activation of JUN/AP-1, TRIF-dependent activation of NF-kappa-B and IRF3, and the phosphorylation of MAPK14/p38. Can restrict West Nile virus (WNV) pathogenesis.[UniProtKB/Swiss-Prot Function]