

Product datasheet for MR217082L3

Cbfa2t2 (NM_172860) Mouse Tagged Lenti ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Cbfa2t2 (NM_172860) Mouse Tagged Lenti ORF Clone
Tag:	Myc-DDK
Symbol:	Cbfa2t2
Synonyms:	A430091M07; C330013D05Rik; Cbfa2t2h; MTGR1
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
E. coli Selection:	Chloramphenicol (34 ug/mL)
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR217082).
Restriction Sites:	SgfI-MluI
Cloning Scheme:	

Cloning sites used for ORF Shuttling:



* The last codon before the Stop codon of the ORF.


ACCN:	NM_172860
ORF Size:	1782 bp




[View online »](#)

OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_172860.2 , NP_766448.1
RefSeq Size:	6100 bp
RefSeq ORF:	1785 bp
Locus ID:	12396
UniProt ID:	O70374
Cytogenetics:	2 H1
Gene Summary:	Transcriptional corepressor which facilitates transcriptional repression via its association with DNA-binding transcription factors and recruitment of other corepressors and histone-modifying enzymes. Via association with PRDM14 is involved in regulation of embryonic stem cell (ESC) pluripotency. Involved in primordial germ cell (PCG) formation (PubMed:27281218). Stabilizes PRDM14 and OCT4 on chromatin in a homooligomerization-dependent mannerCan repress the expression of MMP7 in a ZBTB33-dependent manner (By similarity). Through heteromerization with CBFA2T3/MTG16 may be involved in regulation of the proliferation and the differentiation of erythroid progenitors by repressing the expression of TAL1 target genes (PubMed:19799863). Required for the maintenance of the secretory cell lineage in the small intestine (PubMed:16227606). Can inhibit Notch signaling probably by association with RBPJ and may be involved in GF11-mediated Paneth cell differentiation (PubMed:25398765). [UniProtKB/Swiss-Prot Function]

Product images:



Circular map for MR217082L3



Double digestion of MR217082L3 using SgfI and MluI