

Product datasheet for **MR217131L4V**

Map4k2 (NM_009006) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	Map4k2 (NM_009006) Mouse Tagged ORF Clone Lentiviral Particle
Symbol:	Map4k2
Synonyms:	AI385662; BL44; GCK; Rab8ip
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
Tag:	mGFP
ACCN:	NM_009006
ORF Size:	2463 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR217131).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	NM_009006.2 , NP_033032.1
RefSeq Size:	2466 bp
RefSeq ORF:	2466 bp
Locus ID:	26412
UniProt ID:	Q61161
Cytogenetics:	19 A



[View online »](#)

Gene Summary:

Serine/threonine-protein kinase which acts as an essential component of the MAP kinase signal transduction pathway (PubMed:8643544). Acts as a MAPK kinase kinase (MAP4K) and is an upstream activator of the stress-activated protein kinase/c-Jun N-terminal kinase (SAP/JNK) signaling pathway and to a lesser extent of the p38 MAPKs signaling pathway (By similarity). Required for the efficient activation of JNKs by TRAF6-dependent stimuli, including pathogen-associated molecular patterns (PAMPs) such as polyinosine-polycytidine (poly(IC)), lipopolysaccharides (LPS), lipid A, peptidoglycan (PGN), or bacterial flagellin (By similarity). To a lesser degree, IL-1 and engagement of CD40 also stimulate MAP4K2-mediated JNKs activation (By similarity). The requirement for MAP4K2/GCK is most pronounced for LPS signaling, and extends to LPS stimulation of c-Jun phosphorylation and induction of IL-8 (By similarity). Enhances MAP3K1 oligomerization, which may relieve N-terminal mediated MAP3K1 autoinhibition and lead to activation following autophosphorylation (By similarity). Mediates also the SAP/JNK signaling pathway and the p38 MAPKs signaling pathway through activation of the MAP3Ks MAP3K10/MLK2 and MAP3K11/MLK3 (By similarity). May play a role in the regulation of vesicle targeting or fusion (By similarity).[UniProtKB/Swiss-Prot Function]