

## Product datasheet for **MR217725L3V**

### Syncrip (NM\_019666) Mouse Tagged ORF Clone Lentiviral Particle

#### Product data:

Product Type:	Lentiviral Particles
Product Name:	Syncrip (NM_019666) Mouse Tagged ORF Clone Lentiviral Particle
Symbol:	Syncrip
Synonyms:	2610109K23Rik; 4632417O19Rik; GRY-RBP; hnRNP Q; Nsap1; Nsap1l; pp68
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
Tag:	Myc-DDK
ACCN:	NM_019666
ORF Size:	1869 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR217725).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<a href="#">NM_019666.2</a> , <a href="#">NP_062640.2</a>
RefSeq Size:	2782 bp
RefSeq ORF:	1872 bp
Locus ID:	56403
UniProt ID:	<a href="#">Q7TMK9</a>
Cytogenetics:	9 E3.1



[View online »](#)

**Gene Summary:**

Heterogeneous nuclear ribonucleoprotein (hnRNP) implicated in mRNA processing mechanisms. Component of the CRD-mediated complex that promotes MYC mRNA stability. Isoform 1 and isoform 2 are associated in vitro with pre-mRNA, splicing intermediates and mature mRNA protein complexes. Isoform 1 binds to apoB mRNA AU-rich sequences (By similarity). Isoform 1 is part of the APOB mRNA editosome complex and may modulate the postranscriptional C to U RNA-editing of the APOB mRNA through either by binding to A1CF (APOBEC1 complementation factor), to APOBEC1 or to RNA itself (By similarity). May be involved in translationally coupled mRNA turnover. Implicated with other RNA-binding proteins in the cytoplasmic deadenylation/translational and decay interplay of the FOS mRNA mediated by the major coding-region determinant of instability (mCRD) domain (By similarity). Interacts in vitro preferentially with poly(A) and poly(U) RNA sequences. Isoform 2 may be involved in cytoplasmic vesicle-based mRNA transport through interaction with synaptotagmins.[UniProtKB/Swiss-Prot Function]