

# **Product datasheet for MR218181**

## Rbm8a (NM\_025875) Mouse Tagged ORF Clone

## **Product data:**

**Product Type:** Expression Plasmids

**Product Name:** Rbm8a (NM\_025875) Mouse Tagged ORF Clone

Tag: Myc-DDK
Symbol: Rbm8a

**Synonyms:** 2310057C03Rik; AA673428; Rbm8

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

Cell Selection: Neomycin

ORF Nucleotide >MR218181 representing NM\_025875

Sequence: Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

ATGGCGGACGTGCTGGATCTTCACGAGGCCGGGGGGCGAAGATTTCGCCATGGATGAGGATGGGGACGAAA GCATCCACAAACTAAAAGAAAAAGCAAAGAAACGGAAGGGCCGCGGCTTTGGCTCCGAGGGGTCCCGAGC GCGGATGCGGAGGAGGAGGACGGCGCGGATGAACCTGGACCACAGCGCTCTGTTGAA GGTTGGATTCCTTTGTCACTGGAGTCCACGAAGAAGCCACTGAAGAAGATATCCATGACAAATTCGCTG AATATGGGGAAATAAAAAAATATTCACCTTAATTTGGACAGGCGCACGGGATACTTGAAGGGGTATACTCT AGTTGAATATGAAACATACAAAGAGGCTCAGGCTGCCATGGAAGGACTAAATGGTCAAGATTTGATGGGG CAGCCAATCAGTGTGGACTGGTGTTTTGTTCGTGGACCACCAAAGGGGCAAGAGGAGGAGGAGGACGAAGAC

GAAGCAGGAGTCCAGACCGGAGACGCCGT

**ACGCGT**ACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT

ACAAGGATGACGACGATAAGGTTTAA

Protein Sequence: >MR218181 representing NM\_025875

Red=Cloning site Green=Tags(s)

MADVLDLHEAGGEDFAMDEDGDESIHKLKEKAKKRKGRGFGSEGSRARMREDYDSVEQDGDEPGPQRSVE GWILFVTGVHEEATEEDIHDKFAEYGEIKNIHLNLDRRTGYLKGYTLVEYETYKEAQAAMEGLNGQDLMG

QPISVDWCFVRGPPKGKRRGGRRRSRSPDRRRR

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

**Restriction Sites:** Sgfl-Mlul



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200


CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




#### **Cloning Scheme:**





<sup>\*</sup> The last codon before the Stop codon of the ORF

### Plasmid Map:



**ACCN:** NM\_025875

ORF Size: 519 bp

**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.



Components:

The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
- 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: <u>NM 025875.2</u>, <u>NP 080151.2</u>

 RefSeq Size:
 2628 bp

 RefSeq ORF:
 522 bp

 Locus ID:
 60365

 UniProt ID:
 Q9CWZ3

 Cytogenetics:
 3 F2.1

 MW:
 20.2 kDa

**Gene Summary:** Required for pre-mRNA splicing as component of the spliceosome (By similarity). Core

component of the splicing-dependent multiprotein exon junction complex (EJC) deposited at splice junctions on mRNAs. The EJC is a dynamic structure consisting of core proteins and several peripheral nuclear and cytoplasmic associated factors that join the complex only transiently either during EJC assembly or during subsequent mRNA metabolism. The EJC marks the position of the exon-exon junction in the mature mRNA for the gene expression machinery and the core components remain bound to spliced mRNAs throughout all stages of mRNA metabolism thereby influencing downstream processes including nuclear mRNA export, subcellular mRNA localization, translation efficiency and nonsense-mediated mRNA decay (NMD). Its removal from cytoplasmic mRNAs requires translation initiation from EJC-bearing spliced mRNAs. Associates preferentially with mRNAs produced by splicing. Does not interact with pre-mRNAs, introns, or mRNAs produced from intronless cDNAs. Associates with

both nuclear mRNAs and newly exported cytoplasmic mRNAs (By similarity).

[UniProtKB/Swiss-Prot Function]