

Product datasheet for MR218249L3

Nudt18 (NM_153136) Mouse Tagged Lenti ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Nudt18 (NM_153136) Mouse Tagged Lenti ORF Clone
Tag:	Myc-DDK
Symbol:	Nudt18
Synonyms:	BC036718
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
E. coli Selection:	Chloramphenicol (34 ug/mL)
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR218249).
Restriction Sites:	SgfI-MluI
Cloning Scheme:	

Cloning sites used for ORF Shuttling:



* The last codon before the Stop codon of the ORF.


ACCN:	NM_153136
ORF Size:	969 bp



[View online »](#)

OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_153136.4
RefSeq Size:	3716 bp
RefSeq ORF:	972 bp
Locus ID:	213484
UniProt ID:	Q3U2V3
Cytogenetics:	14 D2
Gene Summary:	Mediates the hydrolysis of oxidized nucleoside diphosphate derivatives. Hydrolyzes 8-oxo-7,8-dihydroguanine (8-oxo-Gua)-containing deoxyribo- and ribonucleoside diphosphates to the monophosphates. Hydrolyzes 8-oxo-dGDP and 8-oxo-GDP with the same efficiencies. Hydrolyzes also 8-OH-dADP and 2-OH-dADP. Exhibited no or minimal hydrolysis activity against 8-oxo-dGTP, 8-oxo-GTP, dGTP, GTP, dGDP and GDP. Probably removes oxidized guanine nucleotides from both the DNA and RNA precursor pools (By similarity). [UniProtKB/Swiss-Prot Function]

Product images:



Circular map for MR218249L3