

## Product datasheet for **MR218479L3V**

### **Rab31 (NM\_133685) Mouse Tagged ORF Clone Lentiviral Particle**

#### **Product data:**

Product Type:	Lentiviral Particles
Product Name:	Rab31 (NM_133685) Mouse Tagged ORF Clone Lentiviral Particle
Symbol:	Rab31
Synonyms:	1700093E07Rik; AI415285; Rab22B
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
Tag:	Myc-DDK
ACCN:	NM_133685
ORF Size:	585 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR218479).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<a href="#">NM_133685.2</a> , <a href="#">NP_598446.2</a>
RefSeq Size:	3476 bp
RefSeq ORF:	588 bp
Locus ID:	106572
UniProt ID:	<a href="#">Q921E2</a>
Cytogenetics:	17 E1.1



[View online »](#)

**Gene Summary:**

The small GTPases Rab are key regulators of intracellular membrane trafficking, from the formation of transport vesicles to their fusion with membranes. Rabs cycle between an inactive GDP-bound form and an active GTP-bound form that is able to recruit to membranes different set of downstream effectors directly responsible for vesicle formation, movement, tethering and fusion. Required for the integrity and for normal function of the Golgi apparatus and the trans-Golgi network. Plays a role in insulin-stimulated translocation of GLUT4 to the cell membrane. Plays a role in the maturation of phagosomes that engulf pathogens, such as S.aureus and Mycobacterium (By similarity). Plays a role in M6PR transport from the trans-Golgi network to endosomes. Plays a role in the internalization of EGFR from the cell membrane into endosomes.[UniProtKB/Swiss-Prot Function]