

Product datasheet for MR218562

Wdr61 (NM_001025376) Mouse Tagged ORF Clone

Product data:

Product Type: Expression Plasmids
Product Name: Wdr61 (NM_001025376) Mouse Tagged ORF Clone
Tag: Myc-DDK
Symbol: Wdr61
Synonyms: 2700038L12Rik; 2810418I05Rik; REC14
Vector: pCMV6-Entry (PS100001)
E. coli Selection: Kanamycin (25 ug/mL)
Cell Selection: Neomycin
ORF Nucleotide Sequence: >MR218562 representing NM_001025376
Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCCGCGATCGCC

ATGACCAACCAAGTACAGTATTCTCTTCAAGCAAGAGCAAGCCCATGATGATGCCATATGGTCAGTTGCCT
GGGAGACAAAACAAAAGGAAAACATTGAAACAGTGGTCACAGGATCCCTGGATGACCTGGTGAAGGTCTG
GAAATGGCGTGATGAGAGGCTGGAGCTCCAGTGGAGCCTGGAGGGACATCAGCTCGGGTGGTGTCTGTG
GACATCAGCCACACTCTCCCATTTGCTGCATCCAGCTCTCTAGACGCTCATATTCGCCTCTGGGACTTGG
AAAATGGCAAACAGATGAAGTCTATAGATGCAGGACCAGTGGATGCCTGGACTTTGGCATTCTCTCTGA
CTCCCAGTATCTGGCCACAGGAACCTCACATGGGAAAGTGAACATTTTTGGTGTGGAAAGTGGAAAAAAA
GAATATTCTTTGGACACTAGAGGAAAATTCATCCTTAGTATTGCATATAGTCCTGATGGGAAATACCTGG
CCAGCGGAGCCATAGACGGAATCATCAATTTTTGACATTGCAACTGGAAAGCTTTTGCATACGCTGGA
AGGCCATGCGATGCCATTTCGCTCCTTGACCTTTCCCTGACTCCCAGCTCCTTGTACGGCTTCAGAT
GATGGCTACATCAAGATCTATGATGTACAACATGCCAATTTGGCTGGCACACTGAGTGCCATGCGTCTC
GGGTGTTGAATGTTGCTTCTGCTGATGACACTCACTTTGTCTCCAGTTCATGACAAAAGGTGAA
GGTTTGGGATGTTGGAACAAGGACCTGTATTCACACCTTCTTTGATCACCAGGATCAGGTTTGGGGAGTA
AAATATAATGAAATGGATCAAAAATTGTATCTGTTGGAGATGACCAGGAAATTCATGTCTATGACTGCC
CAATT

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
ACAAGGATGACGACGATAAGGTTTAA



[View online »](#)

Protein Sequence: >MR218562 representing NM_001025376
 Red=Cloning site Green=Tags(s)

MTNQYSILFKQEQAHDDAIWSVAWETNKKENIETVVTGSLDDLKVKVWKRDERLELQWSLEGHQLGVVSV
 DISHTLPIAASSSLDAHIRLWDLENGKQMKSIDAGPVDATLAFSPDSQYLATGTHMGKVNIFGVESGKK
 EYSLDTRGKFILSIAYSPDGKYLASGAIDGIINIFDIATGKLLHTLEGHAMPIRSLTFSPDSQLLVTASD
 DGYIKIYDVQHANLAGTLSGHASWYLNVAFCPDDTHFVSSSSDKSVKVDVGTRTCIIHTFFDHQDQVWVG
 KYNGNGSKIVSVGDDQEIHYVDCPI


TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Restriction Sites: SgfI-MluI

Cloning Scheme:



Plasmid Map:



ACCN: NM_001025376

ORF Size: 915 bp

OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq Size:	1179 bp
RefSeq ORF:	624 bp
Locus ID:	66317
Cytogenetics:	9 A5.3
MW:	22.9 kDa
Gene Summary:	Component of the PAF1 complex (PAF1C) which has multiple functions during transcription by RNA polymerase II and is implicated in regulation of development and maintenance of embryonic stem cell pluripotency. PAF1C associates with RNA polymerase II through interaction with POLR2A CTD non-phosphorylated and 'Ser-2'- and 'Ser-5'-phosphorylated forms and is involved in transcriptional elongation, acting both independently and synergistically with TCEA1 and in cooperation with the DSIF complex and HTATSF1. PAF1C is required for transcription of Hox and Wnt target genes. PAF1C is involved in hematopoiesis and stimulates transcriptional activity of KMT2A/MLL1. PAF1C is involved in histone modifications such as ubiquitination of histone H2B and methylation on histone H3 'Lys-4' (H3K4me3). PAF1C recruits the RNF20/40 E3 ubiquitin-protein ligase complex and the E2 enzyme UBE2A or UBE2B to chromatin which mediate monoubiquitination of 'Lys-120' of histone H2B (H2BK120ub1); UB2A/B-mediated H2B ubiquitination is proposed to be coupled to transcription. PAF1C is involved in mRNA 3' end formation probably through association with cleavage and poly(A) factors. Required for mono- and trimethylation on histone H3 'Lys-4' (H3K4me3), dimethylation on histone H3 'Lys-79' (H3K4me3). Required for Hox gene transcription. Component of the SKI complex which is thought to be involved in exosome-mediated RNA decay and associates with transcriptionally active genes in a manner dependent on PAF1C (By similarity).[UniProtKB/Swiss-Prot Function]