

Product datasheet for MR218574L3

H2 (H2-Ke6) (NM 013543) Mouse Tagged Lenti ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: H2 (H2-Ke6) (NM_013543) Mouse Tagged Lenti ORF Clone

Tag: Myc-DDK Symbol: H2-Ke6

Synonyms: D17H6S112E; H-2Ke6; Hsd17b8; Ke-6; Ke6; Ring2

Mammalian Cell Puromycin

Selection:

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

E. coli Selection: Chloramphenicol (34 ug/mL)

ORF Nucleotide The ORF insert of this clone is exactly the same as(MR218574).

Sequence:

Restriction Sites: Sgfl-Mlul

Cloning Scheme:





^{*} The last codon before the Stop codon of the ORF.




OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Plasmid Map:



ACCN: NM_013543

ORF Size: 780 bp

OTI Disclaimer: Due to the inherent nature of this plasmid, standard methods to replicate additional amounts

of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at customercom or by

calling 301.340.3188 option 3 for pricing and delivery.

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).



Reconstitution Method:

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 013543.2</u>, <u>NP 038571.2</u>

 RefSeq Size:
 981 bp

 RefSeq ORF:
 780 bp

 Locus ID:
 14979

 UniProt ID:
 P50171

Cytogenetics: 17 17.98 cM

Gene Summary: NAD-dependent 17-beta-hydroxysteroid dehydrogenase with highest activity towards

estradiol. Has very low activity towards testosterone (PubMed:9712896). The heterotetramer with CBR4 has NADH-dependent 3-ketoacyl-acyl carrier protein reductase activity, and thereby plays a role in mitochondrial fatty acid biosynthesis. Within the heterotetramer,

HSD17B8 binds NADH; CBR4 binds NADPD.[UniProtKB/Swiss-Prot Function]