

Product datasheet for MR218596L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Gdf1 (NM_008107) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Gdf1 (NM 008107) Mouse Tagged ORF Clone Lentiviral Particle

Symbol: Gdf1

Synonyms: Al385651; Gdf-1

Mammalian Cell

viaitiitiailati Cei

Puromycin

Selection: Vector:

pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_008107 **ORF Size:** 1074 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(MR218596).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 008107.4, NP 032133.2

 RefSeq Size:
 1419 bp

 RefSeq ORF:
 1074 bp

 Locus ID:
 14559

 UniProt ID:
 P20863

Cytogenetics: 8 B3.3







Gene Summary:

This gene encodes a secreted ligand of the TGF-beta (transforming growth factor-beta) superfamily of proteins. Ligands of this family bind various TGF-beta receptors leading to recruitment and activation of SMAD family transcription factors that regulate gene expression. The encoded preproprotein is proteolytically processed to generate each subunit of the disulfide-linked homodimer. This protein is involved in the establishment of left-right asymmetry in early embryogenesis and in neural development in later embryogenesis. This protein is translated from a monocistronic mRNA early in development, and from a bicistronic mRNA in later stages that also encodes ceramide synthase 1. [provided by RefSeq, Jul 2016]