

## Product datasheet for **MR218937L3V**

### **Puf60 (NM\_133691) Mouse Tagged ORF Clone Lentiviral Particle**

#### **Product data:**

Product Type:	Lentiviral Particles
Product Name:	Puf60 (NM_133691) Mouse Tagged ORF Clone Lentiviral Particle
Symbol:	Puf60
Synonyms:	2410104I19Rik; 2810454F19Rik; SIAHBP1
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
Tag:	Myc-DDK
ACCN:	NM_133691
ORF Size:	1641 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR218937).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<a href="#">NM_133691.5</a> , <a href="#">NP_598452.2</a>
RefSeq Size:	1878 bp
RefSeq ORF:	1644 bp
Locus ID:	67959
UniProt ID:	<a href="#">Q3UEB3</a>
Cytogenetics:	15 D3



[View online »](#)

**Gene Summary:**

DNA- and RNA-binding protein, involved in several nuclear processes such as pre-mRNA splicing, apoptosis and transcription regulation. In association with FUBP1 regulates MYC transcription at the P2 promoter through the core-TFIID basal transcription factor. Acts as a transcriptional repressor through the core-TFIID basal transcription factor. Represses FUBP1-induced transcriptional activation but not basal transcription. Decreases ERCC3 helicase activity. Is also involved in pre-mRNA splicing. Promotes splicing of an intron with weak 3'-splice site and pyrimidine tract in a cooperative manner with U2AF2. Involved in apoptosis induction when overexpressed in HeLa cells. Modulates alternative splicing of several mRNAs. Binds to relaxed DNA of active promoter regions. Binds to the pyrimidine tract and 3'-splice site regions of pre-mRNA; binding is enhanced in presence of U2AF2. Binds to Y5 RNA in association with TROVE2. Binds to poly(U) RNA (By similarity).[UniProtKB/Swiss-Prot Function]