

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for MR219201L4V

Znrf1 (NM_001168622) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	Znrf1 (NM_001168622) Mouse Tagged ORF Clone Lentiviral Particle
Symbol:	Znrf1
Synonyms:	B830022L21Rik; nin283; Rnf42; Zrfp1
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
Tag:	mGFP
ACCN:	NM_001168622
ORF Size:	585 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR219201).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<u>NM 001168622.1</u>
RefSeq Size:	5466 bp
RefSeq ORF:	588 bp
Locus ID:	170737
UniProt ID:	<u>Q91V17</u>
Cytogenetics:	8 E1



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Gene Summary:E3 ubiquitin-protein ligase that mediates the ubiquitination of AKT1 and GLUL, thereby
playing a role in neuron cells differentiation. Plays a role in the establishment and
maintenance of neuronal transmission and plasticity. Regulates Schwann cells differentiation
by mediating ubiquitination of GLUL. Promotes Wallerian degeneration, a neurodegeneration
disorder, by mediating 'Lys-48'-linked polyubiquitination and subsequent degradation of
AKT1 in axons: degradation of AKT1 prevents AKT1-mediated phosphorylation of GSK3B,
leading to GSK3B activation and phosphorylation of DPYSL2/CRMP2 followed by
destabilization of microtubule assembly in axons.[UniProtKB/Swiss-Prot Function]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US