

Product datasheet for **MR219206**

Ffar2 (NM_001168509) Mouse Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Ffar2 (NM_001168509) Mouse Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	Ffar2
Synonyms:	GPCR43; Gpr43
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
ORF Nucleotide Sequence:	>MR219206 ORF sequence Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCC**CGATCGC**

ATGACCCAGACTGGCAGTTCCTTGATCCTCACGGCCTACATCCTCATCTTTCTTACTGGGCTCCCTG
 CCAACCTGCTGGCCCTGCGGGCCTTCATGGGCCGGGTCGCCAGCCTCAGCCTGCCCTGTGCACATCCT
 CCTGCTTAATCTGACCCTGGCGGACTTGCTCCTGTTGCTGCTGCTGCCCTCCGGATCGTGAAGCAGCA
 TCCAACCTCCGCTGGTACCTACCAAAGATCGTGTGCGCGCTGACAGGCTTCGGCTTCTACAGCAGCATCT
 ACTGCAGCACGTGGCTGCTGGCGGGCATCAGCATGGAACGCTACCTGGGAGTGGCCTTCCCGGTGCAGTA
 CAAGTTATCCCGCCGGCCACTGTATGGAGTGATCGCTGCTCTGGTGGCCTGGATCATGTCTTTGGCCAC
 TGCACCATCGTCATCATCGTTACGTACCTGAACTCAACTGAGCAGGTGGGCACTGAGAACCAATAACCT
 GCTACGAGAACTTCACCAAGAGCAGCTGGATGTGGTACTGCCCGTACGACTGGAGCTGTGCCTGGTCTCT
 GTTTTCGTTCCCATGGCAGTCACCATCTTCTGTTATTGGCGCTTCGTGTGGATCATGCTCACGCAGCCC
 CACGTTGGGGCTCAGAGGCGACGCCGGGAGTGGGCTGGCTGTTGTGACGCTTCTTAATTTCTGGTGT
 GCTTTGGACCTACAACATGTCCACCTGGTGGGTTCTACCTGAGGCAGAGCCCTCGTGGCGGGTGA
 GGCTGTGGTGTTCAGTTCCCTCAATGCCAGCCTGGATCCATTATTGTTCTACTTCTCCTCCTCCGTGGT
 CGCAGAGCTTTTGGAAAGGTTTGCTACTGATCCGCAATCCTGCCTCCTATGCTGGGCAGGGGAGCCA
 AAGAGACAGTGGAGGGGACCAAGATGGACAGGGGTGGAAGTCAAGCAGAAGGGGTACAGAGTTCTGAATT
 TGTCACCGAG

ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
 ACAAGGATGACGACGATAAGGTTTAA


[View online »](#)

Protein Sequence: >MR219206 protein sequence
Red=Cloning site Green=Tags(s)



MTPDWHSSLLITAYILIFLTGLPANLLALRAFMGRVRQPQAPVHILLNLTADLLLLLLLPRIVEAA
SNFRWYLPKIVCALTGFIFYSSYCYSTWLLAGISMERYLGVAFPVQYKLSRRPLYGVIAALVAVIMSF
GHCITIVIIVQYLRNSTEQVTGENQITCYENFTQEQLDVVLPVRLLELVLVFFVPMAVTIFCYWRFVWIMLTQ
P
HVGAGRRRRRAVLGAVVLTLLNLVCFGPYNNMSHLVGFYLRQSPSRWRVEAVFVSSLNASLDPLLFYFSSSV
RRAFGKGLLLIRNPASSMLGRGAKETVEGTGKMDRGQSAEGVOSFEVTE

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Restriction Sites: Sgfl-MluI

Cloning Scheme:

Cloning sites used for ORF Shuttling:



* The last codon before the Stop codon of the ORF

ACCN: NM 001168509

ORF Size: 990 bp


OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:	<ol style="list-style-type: none"> 1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
Note:	Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.
RefSeq:	<u>NM_001168509.1, NP_001161981.1</u>
RefSeq Size:	1969 bp
RefSeq ORF:	993 bp
Locus ID:	233079
UniProt ID:	<u>Q8VCK6</u>
Cytogenetics:	7 B1
MW:	37.1 kDa
Gene Summary:	<p>G protein-coupled receptor that is activated by a major product of dietary fiber digestion, the short chain fatty acids (SCFAs), and that plays a role in the regulation of whole-body energy homeostasis and in intestinal immunity. In omnivorous mammals, the short chain fatty acids acetate, propionate and butyrate are produced primarily by the gut microbiome that metabolizes dietary fibers. SCFAs serve as a source of energy but also act as signaling molecules. That G protein-coupled receptor is probably coupled to the pertussis toxin-sensitive, G(i/o)-alpha family of G proteins but also to the Gq family (PubMed:23589301). Its activation results in the formation of inositol 1,4,5-trisphosphate, the mobilization of intracellular calcium, the phosphorylation of the MAPK3/ERK1 and MAPK1/ERK2 kinases and the inhibition of intracellular cAMP accumulation. May play a role in glucose homeostasis by regulating the secretion of GLP-1, in response to short-chain fatty acids accumulating in the intestine (PubMed:22190648, PubMed:23589301). May also regulate the production of LEP/Leptin, a hormone acting on the central nervous system to inhibit food intake (PubMed:20399779). Finally, may also regulate whole-body energy homeostasis through adipogenesis regulating both differentiation and lipid storage of adipocytes (PubMed:16123168, PubMed:23589301). In parallel to its role in energy homeostasis, may also mediate the activation of the inflammatory and immune responses by SCFA in the intestine, regulating the rapid production of chemokines and cytokines (PubMed:23665276). May also play a role in the resolution of the inflammatory response and control chemotaxis in neutrophils (PubMed:19917676, PubMed:19865172). In addition to SCFAs, may also be activated by the extracellular lectin FCN1 in a process leading to activation of monocytes and inducing the secretion of interleukin-8/IL-8 in response to the presence of microbes. [UniProtKB/Swiss-Prot Function]</p>

Product images:



Circular map for MR219206