

## Product datasheet for **MR219389L3V**

### Chst11 (NM\_021439) Mouse Tagged ORF Clone Lentiviral Particle

#### Product data:

|                           |  |
|---------------------------|--|
| Product Type:             | Lentiviral Particles   |
| Product Name:             | Chst11 (NM_021439) Mouse Tagged ORF Clone Lentiviral Particle  |
| Symbol:                   | Chst11   |
| Synonyms:                 | 1110020P09Rik; C4s; C4ST; C4ST-1; C4ST1  |
| Mammalian Cell Selection: | Puromycin  |
| Vector:                   | pLenti-C-Myc-DDK-P2A-Puro (PS100092)   |
| Tag:                      | Myc-DDK  |
| ACCN:                     | NM_021439  |
| ORF Size:                 | 1056 bp  |
| ORF Nucleotide Sequence:  | The ORF insert of this clone is exactly the same as(MR219389).   |
| OTI Disclaimer:           | The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a> |
| OTI Annotation:           | This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.   |
| RefSeq:                   | <a href="#">NM_021439.2</a> , <a href="#">NP_067414.2</a>  |
| RefSeq Size:              | 5532 bp  |
| RefSeq ORF:               | 1059 bp  |
| Locus ID:                 | 58250  |
| UniProt ID:               | <a href="#">Q9JME2</a>   |
| Cytogenetics:             | 10 C1  |



[View online »](#)

**Gene Summary:**

Catalyzes the transfer of sulfate to position 4 of the N-acetylgalactosamine (GalNAc) residue of chondroitin. Chondroitin sulfate constitutes the predominant proteoglycan present in cartilage and is distributed on the surfaces of many cells and extracellular matrices. Can also sulfate Gal residues in desulfated dermatan sulfate. Preferentially sulfates in GlcA->GalNAc unit than in IdoA->GalNAc unit. Does not form 4, 6-di-O-sulfated GalNAc when chondroitin sulfate C is used as an acceptor.[UniProtKB/Swiss-Prot Function]