

Product datasheet for MR219506L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Tead4 (NM 011567) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Tead4 (NM_011567) Mouse Tagged ORF Clone Lentiviral Particle

Symbol: Tead4

Synonyms: Etfr; ETFR-; ETFR-2; Etfr2; FR-19; Rtef; Rtef1; Tcf13r1; Tcf13r1; TEAD-4; Tef; TEF-3; Tef3;

Tefr; Tefr1; Tefr1a

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_011567 **ORF Size:** 1361 bp

ORF Nucleotide

Sequence:

The ORF insert of this clone is exactly the same as(MR219506).

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 011567.2</u>

 RefSeq Size:
 3076 bp

 RefSeq ORF:
 1284 bp

 Locus ID:
 21679

 UniProt ID:
 Q62296

 Cytogenetics:
 6 62.92 cM







Gene Summary:

This gene product is a member of the transcriptional enhancer factor (TEF) family of transcription factors, which contain the TEA/ATTS DNA-binding domain. It is preferentially expressed in the skeletal muscle, and binds to the M-CAT regulatory element found in promoters of muscle-specific genes to direct their gene expression. This factor may play a role in the embryonic development of skeletal muscle. Alternatively spliced transcripts encoding distinct isoforms, which are translated through the use of a non-AUG (AUU) initiation codon, have been described for this gene. [provided by RefSeq, Jul 2008]