

Product datasheet for MR219511L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Nrcam (NM_176930) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Nrcam (NM_176930) Mouse Tagged ORF Clone Lentiviral Particle

Symbol: Nrcam

Synonyms: Bravo; C030017F07Rik; C130076O07Rik; mKIAA0343

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_176930 **ORF Size:** 3768 bp

ORF Nucleotide

TI 005 ' ' C '

Sequence:

The ORF insert of this clone is exactly the same as(MR219511).

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This

clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 176930.4, NP 795904.3

 RefSeq Size:
 7572 bp

 RefSeq ORF:
 3771 bp

 Locus ID:
 319504

 UniProt ID:
 <u>Q810U4</u>

Cytogenetics: 12 20.71 cM







Gene Summary:

Cell adhesion protein that is required for normal responses to cell-cell contacts in brain and in the peripheral nervous system. Plays a role in neurite outgrowth in response to contactin binding (PubMed:11564762). Plays a role in mediating cell-cell contacts between Schwann cells and axons (PubMed:20188654). Plays a role in the formation and maintenance of the nodes of Ranvier on myelinated axons. Nodes of Ranvier contain clustered sodium channels that are crucial for the saltatory propagation of action potentials along myelinated axons. During development, nodes of Ranvier are formed by the fusion of two heminodes. Required for normal clustering of sodium channels at heminodes; not required for the formation of mature nodes with normal sodium channel clusters (PubMed:14602817, PubMed:20188654). Required, together with GLDN, for maintaining NFASC and sodium channel clusters at mature nodes of Ranvier (PubMed:24719088).[UniProtKB/Swiss-Prot Function]