

Product datasheet for MR219719L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Hist1h2be (H2bc6) (NM_001177653) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Hist1h2be (H2bc6) (NM_001177653) Mouse Tagged ORF Clone Lentiviral Particle

Symbol: H2bc6

Synonyms: Gm11398; H2bc4; H2bc8; Hist1h2; Hist1h2be

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_001177653

ORF Size: 378 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(MR219719).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: NM 001177653.1, NP 001171124.1

 RefSeq Size:
 2483 bp

 RefSeq ORF:
 381 bp

 Locus ID:
 319179

 UniProt ID:
 Q6ZWY9

 Cytogenetics:
 13 A3.1





Gene Summary:

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chromatin into higher order structures. This gene encodes a replication-dependent histone that is a member of the histone H2B family and generates multiple transcripts through alternative splicing, the use of the conserved stem-loop termination motif, and the polyA addition motif. [provided by RefSeq, Aug 2015]