


OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_172260.3 , NP_758464.2
RefSeq Size:	4810 bp
RefSeq ORF:	2202 bp
Locus ID:	216543
UniProt ID:	Q8C0D9
Cytogenetics:	11 12.92 cM
Gene Summary:	Involved in maintenance of centrosome cohesion, probably as part of a linker structure which prevents centrosome splitting. Required for localization of CDK5RAP2 to the centrosome during interphase.[UniProtKB/Swiss-Prot Function]

Product images:



Circular map for MR219870L3