

Product datasheet for MR219928L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Exd1 (NM_172857) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Exd1 (NM 172857) Mouse Tagged ORF Clone Lentiviral Particle

Symbol: Exd1

Synonyms: 4932702D22Rik; Exdl1; mExd1

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

 Tag:
 Myc-DDK

 ACCN:
 NM_172857

ORF Size: 1710 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(MR219928).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 172857.2</u>, <u>NP 766445.1</u>

 RefSeq Size:
 3143 bp

 RefSeq ORF:
 1713 bp

 Locus ID:
 241624

 UniProt ID:
 Q8CDF7

 Cytogenetics:
 2 E5



Gene Summary:

RNA-binding component of the PET complex, a multiprotein complex required for the processing of piRNAs during spermatogenesis. The piRNA metabolic process mediates the repression of transposable elements during meiosis by forming complexes composed of piRNAs and Piwi proteins and governs the methylation and subsequent repression of transposable elements, preventing their mobilization, which is essential for the germline integrity (PubMed:26669262). The PET complex is required during the secondary piRNAs metabolic process for the PIWIL2 slicing-triggered loading of PIWIL4 piRNAs. In the PET complex, EXD1 probably acts as an RNA adapter. EXD1 is an inactive exonuclease (By similarity).[UniProtKB/Swiss-Prot Function]