

Product datasheet for MR220358L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Arfgap1 (NM_145760) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Arfgap1 (NM_145760) Mouse Tagged ORF Clone Lentiviral Particle

Symbol: Arfgap1

Synonyms: Al115377; Arf1gap

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag:Myc-DDKACCN:NM_145760

ORF Size: 1242 bp

ORF Nucleotide Sequence:

The ORF insert of this clone is exactly the same as(MR220358).

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This

clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 145760.3, NP 665703.2

 RefSeq Size:
 2721 bp

 RefSeq ORF:
 1245 bp

 Locus ID:
 228998

 UniProt ID:
 Q9EPJ9

Cytogenetics: 2 103.53 cM







Gene Summary:

GTPase-activating protein (GAP) for the ADP ribosylation factor 1 (ARF1). Involved in membrane trafficking and /or vesicle transport. Promotes hydrolysis of the ARF1-bound GTP and thus, is required for the dissociation of coat proteins from Golgi-derived membranes and vesicles, a prerequisite for vesicle's fusion with target compartment. Probably regulates ARF1-mediated transport via its interaction with the KDELR proteins and TMED2. Overexpression induces the redistribution of the entire Golgi complex to the endoplasmic reticulum, as when ARF1 is deactivated. Its activity is stimulated by phosphoinosides and inhibited by phosphatidylcholine (By similarity).[UniProtKB/Swiss-Prot Function]